



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C07D 307/87, A61K 31/343</b>		<b>A1</b>	(11) International Publication Number: <b>WO 00/34263</b>
			(43) International Publication Date: <b>15 June 2000 (15.06.00)</b>
(21) International Application Number: <b>PCT/DK99/00676</b>		(74) Common Representative: <b>H. LUNDBECK A/S; PETERSEN, John, Meidahl, Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).</b>	
(22) International Filing Date: <b>3 December 1999 (03.12.99)</b>			
(30) Priority Data: 60/111,360 8 December 1998 (08.12.98) US PA 1998 01631 9 December 1998 (09.12.98) DK		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): <b>H. LUNDBECK A/S [DK/DK]; Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).</b>			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): <b>ANDERSEN, Kim [DK/DK]; Ringerbakken 22, DK-2830 Virum (DK). RÖTTLÄNDER, Mario [DE/DK]; Harrestrupvang 3c, 2th, DK-2500 Valby (DK). BØGESØ, Klaus, Peter [DK/DK]; Hørsholm Park 16, 2tv, DK-2970 Hørsholm (DK). PEDERSEN, Henrik [DK/DK]; Mellemvangen 63, DK-2700 Brønshøj (DK). RUHLAND, Thomas [DE/DK]; Østergårds Alle 16, DK-2500 Valby (DK). DANCER, Robert [AU/DK]; J.M. Thieles Vej 8, st th, DK-1961 Frederiksberg C (DK).</b>		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: <b>BENZOFURAN DERIVATIVES, THEIR PREPARATION AND USE</b>			
<div style="text-align: center;">( I )</div>			
<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  ( 1 ) </div> <div style="text-align: center;">  ( 2 ) </div> <div style="text-align: center;">  ( 3 ) </div> <div style="text-align: center;">  ( 4 ) </div> </div>			
(57) Abstract			
<p>The present invention relates to benzofuran derivatives having general Formula (I). A is selected from (1), (2), (3), (4) wherein: Z is O or S; s is 0 or 1; q is 0 or 1; R<sup>4</sup> is hydrogen, C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl, C<sub>2-6</sub>-alkynyl, C<sub>1-6</sub>-alkyl-Aryl, or C<sub>1-6</sub>-alkyl-O-Aryl; D is a spacer group selected from branched or straight chain C<sub>1-6</sub>-alkylene, C<sub>2-6</sub>-alkenylene and C<sub>2-6</sub>-alkynylene; its enantiomers, and pharmaceutically acceptable acid addition salt thereof. The compounds are potently binding to the 5-HT<sub>1A</sub> receptor.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**Benz furan derivatives, their preparation and use**

The present invention relates to novel benzofuran derivatives potently binding to the 5-HT<sub>1A</sub> receptor, pharmaceutical compositions containing these compounds and the use thereof for the treatment of certain psychiatric and neurological disorders. Many of the compounds of the invention are also potent serotonin reuptake inhibitors and are considered to be particularly useful for the treatment of depression.

**Background Art**

10

Clinical studies of known 5-HT<sub>1A</sub> partial agonists such as e.g. buspirone, ipsapirone and gepirone have shown that 5-HT<sub>1A</sub> partial agonists are useful in the treatment of anxiety disorders such as generalised anxiety disorder, panic disorder, and obsessive compulsive disorder (Glitz, D. A., Pohl, R., *Drugs* 1991, 41, 11). Preclinical studies indicate that full agonists also are useful in the treatment of the above mentioned anxiety related disorders (Schipper, *Human Psychopharm.*, 1991, 6, S53).

There is also evidence, both clinical and preclinical, in support of a beneficial effect of 5-HT<sub>1A</sub> partial agonists in the treatment of depression as well as impulse control disorders and alcohol abuse (van Hest, *Psychopharm.*, 1992, 107, 474; Schipper et al, *Human Psychopharm.*, 1991, 6, S53; Cervo et al, *Eur. J. Pharm.*, 1988, 158, 53; Glitz, D. A., Pohl, R., *Drugs* 1991, 41, 11; Grof et al., *Int. Clin. Psychopharmacol.* 1993, 8, 167-172; Anseau et al., *Human Psychopharmacol.* 1993, 8, 279-283).

5-HT<sub>1A</sub> agonists and partial agonists inhibit isolation-induced aggression in male mice indicating that these compounds are useful in the treatment of aggression (Sánchez et al, *Psychopharmacology*, 1993, 110, 53-59).

Furthermore, 5-HT<sub>1A</sub> agonists have been reported to show activity in animal models predictive for antipsychotic effects (Wadenberg and Ahlenius, *J. Neural. Transm.*, 1991, 83, 43; Ahlenius, *Pharmacol. & Toxicol.*, 1989, 64, 3; Lowe et al., *J. Med. Chem.*, 1991, 34, 1860; New et al., *J. Med. Chem.*, 1989, 32, 1147; and Martin et al., *J. Med. Chem.*,

1989, 32, 1052) and may therefore be useful in the treatment of psychotic disorders such as schizophrenia. Recent studies also indicate that 5-HT<sub>1A</sub> receptors are important in the serotonergic modulation of haloperidol-induced catalepsy (Hicks, *Life Science* 1990, 47, 1609) suggesting that 5-HT<sub>1A</sub> agonists are useful in the treatment of the side effects  
5 induced by conventional antipsychotic agents such as e.g. haloperidol.

5-HT<sub>1A</sub> agonists have shown neuroprotective properties in rodent models of focal and global cerebral ischaemia and may, therefore, be useful in the treatment of ischaemic disease states (Prehn, *Eur. J. Pharm.* 1991, 203, 213).

10

Pharmacological studies have been presented which indicate that 5-HT<sub>1A</sub> antagonists are useful in the treatment of senile dementia (Bowen et al., *Trends Neur. Sci.* 1992, 15, 84).

An overview of 5-HT<sub>1A</sub> antagonists and proposed potential therapeutic targets for these  
15 antagonists based upon preclinical and clinical data are presented by Schechter et al., *Serotonin*, 1997, Vol.2, Issue 7. It is stated that 5-HT<sub>1A</sub> antagonists may be useful in the treatment of schizophrenia, dementia associated with Alzheimer's disease, and in combination with SSRI antidepressants also to be useful in the treatment of depression.

20 Both in animal models and in clinical trials it has been shown that 5-HT<sub>1A</sub> agonists exert antihypertensive effects via a central mechanism (Saxena and Villalón, *Trends Pharm. Sci.* 1990, 11, 95; Gillis et al., *J. Pharm. Exp. Ther.* 1989, 248, 851). 5-HT<sub>1A</sub> ligands may, therefore, be beneficial in the treatment of cardiovascular disorders.

25 5-HT reuptake inhibitors are well known antidepressant drugs and useful for the treatment of panic disorders and social phobia.

The effect of combined administration of a compound that inhibits serotonin reuptake and a 5-HT<sub>1A</sub> receptor antagonist has been evaluated in several studies (Innis, R.B. et al., *Eur. J. Pharmacol.*, 1987, 143, p 195-204 and Gartside, S.E., *Br. J. Pharmacol.* 1995, 115, p  
30 1064-1070, Blier, P. et al, *Trends Pharmacol. Sci.* 1994, 15, 220). In these studies it was found that 5-HT<sub>1A</sub> receptor antagonists would abolish the initial brake on 5-HT

neurotransmission induced by the serotonin reuptake inhibitors and thus produce an immediate boost of 5-HT transmission and a more rapid onset of therapeutic action.


Several patent applications have been filed which cover the use of a combination of a 5-HT<sub>1A</sub> antagonist and a serotonin reuptake inhibitor for the treatment of depression (see EP-A2-687 472 and EP-A2-714 663).

Accordingly, agents acting on the 5-HT<sub>1A</sub> receptor, both agonists and antagonists, are believed to be of potential use in the therapy of psychiatric and neurological disorders and thus being highly desired. Furthermore, antagonists at the same time having potent serotonin reuptake inhibition activity may be useful for the treatment of depression.

### Summary of the Invention

It has now been found that compounds of a certain class of benzofuran derivatives bind to the 5-HT<sub>1A</sub> receptor with high affinities. Furthermore, it has been found that many of these compounds have potent serotonin reuptake inhibition activity.

Accordingly, the present invention relates to novel compounds of the general Formula I:



wherein





R¹ is hydrogen, halogen, trifluoromethyl, trifluoromethylsulfonyloxy, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-8</sub> cycloalkyl, C<sub>1-6</sub> alkoxy, hydroxy, formyl, acyl, amino, C<sub>1-6</sub> alkylamino, C<sub>2-12</sub> dialkylamino, acylamino, C<sub>1-6</sub> alkoxycarbonylamino, aminocarbonylamino, C<sub>1-6</sub> alkylaminocarbonylamino, C<sub>2-12</sub> dialkylaminocarbonylamino, nitro, cyano, COOH, or COO-C<sub>1-6</sub> alkyl;

$R^2$  and  $R^3$  are each independently selected from hydrogen, trifluoromethyl,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl,  $C_{3-8}$  cycloalkyl and  $C_{1-6}$  alkoxy;

$n$  is 1, 2, 3, 4 or 5;

5  $m$  is 0 or 1;

A is selected from the following groups:



wherein

10  $Z$  is O or S;

$s$  is 0 or 1;




$q$  is 0 or 1;

$R^4$  is hydrogen,  $C_{1-6}$ -alkyl,  $C_{2-6}$ -alkenyl,  $C_{2-6}$ -alkynyl,  $C_{1-6}$ -alkyl-Aryl, or  $C_{1-6}$ -alkyl-O-Aryl,

15

D is a spacer group selected from branched or straight chain  $C_{1-6}$ -alkylene,  $C_{2-6}$ -alkenylene and  $C_{2-6}$ -alkynylene;

B is a group selected from a group of formula (II), (III), and (IV)



20 wherein  $R^5$ ,  $R^6$ ,  $R^7$ ,  $R^8$ ,  $R^9$  and  $R^{10}$  are each independently selected among the  $R^1$  substituents;

or  $R^8$  and  $R^9$  together form a fused 5- or 6-membered ring optionally containing further heteroatoms;

or two of the groups of  $R^5$ ,  $R^6$  and  $R^7$  are linked together thereby forming a

$\text{—O—(CH}_2\text{)}_p\text{—O—}$  -bridge wherein p is 1 or 2;

- 5 Ar and Aryl are independently selected from the group consisting of phenyl, 2-thienyl, 3-thienyl, 2-furanyl, 3-furanyl, 2-pyrimidyl, 1-indolyl, 2-indolyl, 3-indolyl, indol-2-on-1-yl, indol-2-on-3-yl, 2- or 3-benzofuranyl, 2- or 3-benzothiophenyl, 1-naphthyl or 2-naphthyl, each optionally substituted with halogen,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy,  $C_{1-6}$  alkylthio, hydroxy,  $C_{1-6}$  alkylsulfonyl, cyano, trifluoromethyl, trifluoromethylsulfonyloxy,  $C_{3-8}$
- 10 cycloalkyl,  $C_{3-8}$  cycloalkyl- $C_{1-6}$  alkyl, nitro, amino,  $C_{1-6}$  alkylamino,  $C_{2-12}$  dialkylamino, acylamino or alkylenedioxy;

its enantiomers, and pharmaceutically acceptable acid addition salt thereof.

- 15 In one embodiment of the invention A is a group of formula (1) and the other substituents are as defined above.

In another embodiment of the invention A is a group of formula (2) and the other substituents are as defined above.

20

In a third embodiment of the invention A is a group of formula (3) and the other substituents are as defined above.

- 25 In a fourth embodiment of the invention A is a group of formula (4) and the other substituents are as defined above.

Thus in a preferred embodiment of the invention A is a group of formula (1) and  $R^4$  is methyl, ethyl, propyl, prop-2-en-1-yl, 2-furylmethyl, or 2-phenoxyethyl;  $q = 0$ ; or A is a group of formula (1) and Z is O and the other substituents are as defined above.

30

In a further embodiment of the invention, B is a group of formula (II), preferably a alkoxy-substituted phenyl, a benzodioxan group or a 1,2-methylenedioxybenzene group and the other substituents are as defined above.

In a further embodiment of the invention, B is a group of formula (III), preferably a 3-indolyl group and the other substituents are as defined above.

In a further embodiment of the invention, B is a group of formula (III), preferably a 3-indolyl group and the substituents R<sup>8</sup> and R<sup>9</sup> are preferably selected from hydrogen, methyl, fluoro, chloro, bromo, iodo, *t*-butyl or *i*-propyl in the 5-position; or fluoro, chloro or carboxy in the 7-position; or by 5,7-difluoro, 4-fluoro-7-methyl or 4-chloro-7-methyl; or the two substituents together form a pyridyl ring fused to the 3-indolyl.

In a further embodiment of the invention, B is a group of formula (IV) and the other substituents are as defined above.

Ar is preferably phenyl or phenyl substituted with halogen or CF<sub>3</sub>, most preferably substituted with F or Cl in the 4-position or Cl or CF<sub>3</sub> in the 3-position.

R<sup>1</sup> is preferably H, CN or F in the 5-position of the isobenzofuran group.

R<sup>2</sup> and R<sup>3</sup> are preferably selected from hydrogen or methyl.

n is preferably 2, 3 or 4.

m is preferably 0.

In a preferred embodiment of the invention n = 2, 3 or 4; R<sup>2</sup> and R<sup>3</sup> are both hydrogen; R<sup>1</sup> is H, CN or F in the 5-position of the isobenzofuran group; and Ar is phenyl which may be substituted with F or Cl in the 4-position or with Cl or CF<sub>3</sub> in the 3-position and the other substituents are as defined above.



In another preferred embodiment of the invention, A is a group of formula (1);  $q = 0$ ;  $R^4$  is methyl; D is propylene;  $m = 0$ ; and B is a 1,4-benzodioxan group of Formula (II) attached in the 5-position and the other substituents are as defined above.

- 5 In another preferred embodiment of the invention, A is a group of formula (1);  $R^4$  is  $CH_3$  or prop-2-en-1-yl;  $n = 3$ ; D is ethylene or propylene; and B is a phenyl group wherein at least one substituent is OMe and the other substituents are as defined above.

- 10 In a further embodiment of the invention, A is a group of formula (1);  $q$  is 0;  $R^4$  is methyl, ethyl, propyl, 2-propen-1-yl, 2-furylmethyl or 2-phenoxyethyl; D is ethylene, propylene or butylene;  $m = 0$ ; and B is a 3-indolyl group of Formula (III) and the other substituents are as defined above.

- 15 In another preferred embodiment of the invention, A is a group of formula (2) or (3);  $n = 3$ ;  $m = 0$ ; and B is an 4- or 5-indolyl-group of Formula (IV) wherein  $R^{10}$  is hydrogen;  $R^1$  is CN in the 5-position of the isobenzofuran and Ar is 4-Fluorophenyl and the other substituents are as defined above.

- 20 The invention also relates to a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof and at least one pharmaceutically acceptable carrier or diluent.

- 25 In a further embodiment, the invention relates to the use of a compound of formula (I) or a pharmaceutically acceptable acid addition salt thereof for the preparation of a medicament for the treatment of a disorder or disease responsive to the effect of  $5-HT_{1A}$  receptors.

- 30 In particular, the invention relates to the use of a compound according to the invention or a pharmaceutically acceptable acid addition salt thereof for the preparation of a medicament for the treatment of depression, psychosis, anxiety disorders, panic disorder, obsessive compulsive disorder, impulse control disorder, alcohol abuse, aggression, ischaemia, senile dementia, cardiovascular disorders or social phobia.

In still another embodiment, the present invention relates to a method for the treatment of a disorder or disease of living animal body, including a human, which is responsive to the effect of 5-HT<sub>1A</sub> receptors comprising administering to such a living animal body, including a human, a therapeutically effective amount of a compound of formula (I) or a  
5 pharmaceutically acceptable acid addition salt thereof.

The compounds of the invention have high affinity for the 5-HT<sub>1A</sub> receptor. Accordingly, the compounds of the invention are considered useful for the treatment of depression, psychosis, anxiety disorders, such as generalised anxiety disorder, panic disorder, and  
10 obsessive compulsive disorder, impulse control disorder, alcohol abuse, aggression, ischaemia, senile dementia, cardiovascular disorders and social phobia.

Due to their combined antagonism of 5-HT<sub>1A</sub> receptors and serotonin reuptake inhibiting effect, many of the compounds of the invention are considered particularly useful as fast  
15 onset of action medicaments for the treatment of depression. The compounds may also be useful for the treatment of depression in patients who are resistant to treatment with currently available antidepressants.

#### Detailed Description of the Invention

20 Some of the compounds of general Formula I may exist as optical isomers thereof and such optical isomers are also embraced by the invention.

The term C<sub>1-6</sub> alkyl refers to a branched or unbranched alkyl group having from one to six carbon atoms inclusive, such as methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-  
25 methyl-2-propyl and 2-methyl-1-propyl.

Similarly, C<sub>2-6</sub> alkenyl and C<sub>2-6</sub> alkynyl, respectively, designate such groups having from two to six carbon atoms, inclusive.

30 Halogen means fluoro, chloro, bromo, or iodo.

The term C<sub>3,8</sub> cycloalkyl designates a monocyclic or bicyclic carbocycle having three to eight C-atoms, such as cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

The terms C<sub>1,6</sub> alkoxy, C<sub>1,6</sub> alkylthio, C<sub>1,6</sub> alkylsulfonyl, designate such groups in which the  
5 alkyl group is C<sub>1,6</sub> alkyl as defined above.

Acyl means -CO-alkyl wherein the alkyl group is C<sub>1,6</sub> alkyl as defined above.

C<sub>1,6</sub> alkylamino means -NH-alkyl, and C<sub>2,12</sub> dialkylamino means -N-(alkyl)<sub>2</sub> where the alkyl  
10 group is C<sub>1,6</sub> alkyl as defined above.

Acylamino means -NH-acyl wherein acyl is as defined above.

C<sub>1,6</sub> alkoxycarbonylamino means alkyl-O-CO-NH- wherein the alkyl group is C<sub>1,6</sub> alkyl as  
15 defined above.

C<sub>1,6</sub> alkylaminocarbonylamino means alkyl-NH-CO-NH- wherein the alkyl group is  
C<sub>1,6</sub> alkyl as defined above.

20 C<sub>2,12</sub> dialkylaminocarbonylamino means (alkyl)<sub>2</sub>-N-CO-NH- wherein the alkyl group is  
C<sub>1,6</sub> alkyl as defined above.

Exemplary of organic acid addition salts according to the invention are those with maleic, fumaric, benzoic, ascorbic, succinic, oxalic, bis-methylenesalicylic, methanesulfonic,  
25 ethanedisulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, and theophylline acetic acids, as well as the 8-halotheophyllines, for example 8-bromotheophylline. Exemplary of inorganic acid addition salts according to the invention are those with hydrochloric, hydrobromic, sulfuric,  
30 sulfamic, phosphoric, and nitric acids. The acid addition salts of the invention are preferably pharmaceutically acceptable salts formed with non-toxic acids.

Furthermore, the compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention.

5

Some of the compounds of the present invention contain chiral centres and such compounds exist in the form of isomers (e.g. enantiomers). The invention includes all such isomers and any mixtures thereof including racemic mixtures.


- 10 Racemic forms can be resolved into the optical antipodes by known methods, for example, by separation of diastereomeric salts thereof with an optically active acid, and liberating the optically active amine compound by treatment with a base. Another method for resolving racemates into the optical antipodes is based upon chromatography on an optically active matrix. Racemic compounds of the present invention can thus be resolved
- 15 into their optical antipodes, e.g., by fractional crystallisation of d- or l- (tartrates, mandelates, or camphorsulphonate) salts for example. The compounds of the present invention may also be resolved by the formation of diastereomeric derivatives.

- Additional methods for the resolution of optical isomers, known to those skilled in the art,
- 20 may be used. Such methods include those discussed by J. Jaques, A. Collet, and S. Wilen in "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981).

Optically active compounds can also be prepared from optically active starting materials.


- 25 The compounds of the invention can be prepared by one of the following methods comprising:

a) alkylating an amine of formula




wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $n$  and  $Ar$  are as defined above with an alkylating agent of formula  $G-(D)_s-(Z)_q-(CH_2)_m-B$  wherein  $D$ ,  $Z$ ,  $m$ ,  $s$ ,  $q$  and  $B$  are as defined above and  $G$  is a suitable  
 5 leaving group such as halogen, mesylate, or tosylate;

b) alkylating an amine of formula  $H-A-(CH_2)_m-B$  wherein  $A$ ,  $m$  and  $B$  are as defined above with an alkylating agent of formula




wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $n$  and  $Ar$  are as defined above and  $G$  is a suitable leaving group such as halogen, mesylate, or tosylate;

c) reductive alkylation of an amine of formula



wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $n$  and  $Ar$  are as defined above with an aldehyde of formula




(VIII)

wherein Z, m, q and B are as defined above and t is 1-5;

5


d) reducing an amide of formula



(XI)

10 wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, n, q, Ar, Z, m and B are as defined above and t is 1-5;

e) releasing final product by the means of Hofmann elimination from a resin of formula




15

(XII)

wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, n, s, q, Ar, D, Z, m and B are as defined above, G is as defined above; and HOR' is a hydroxy substituted resin such as cross linked hydroxymethylpolystyrene or Wang resin.


13

f) reacting a compound of the formula



XIII

wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ , Ar, D and N are as defined above;  $(OH)_2Q$  is a diol such as substituted ethylene glycol or propylene glycol, or a polymer bound diol, with a hydrazine  
 5 of formula



XIV

wherein  $R^8$  and  $R^9$  is as defined above, using Lewis acids as catalyst.

10

The alkylations according to Methods a and b are generally performed by boiling the reactants under reflux or by heating them at a fixed temperature in a suitable solvent such as acetone, methyl isobutyl ketone, tetrahydrofuran, dioxane, ethanol, 2-propanol, ethyl acetate, N,N-dimethylformamide, dimethyl sulfoxide or 1-methyl-2-pyrrolidinone in the  
 15 presence of a base such as triethylamine or potassium carbonate. Amines of formula V are prepared by means of demethylation according to the method described by Bigler et al, Eur. J. Med. Chem. Chim. Ther, 1977, 12, 289-295, or by the methods outlined in examples 14 and 15. The starting materials used in example 14 were prepared as described in example 9 or from readily available compounds by standard methods. The enantiomers  
 20 of 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile used as starting material for the demethylation are prepared as described in EP patent No. 347066. The alkylating agents of formula  $G-(D)_s-(Z)_q-(CH_2)_m-B$  are

commercially available, prepared by methods obvious to the chemist skilled in the art or prepared as exemplified in Examples 5-8. Ethyl 1,4-benzodioxan-5-carboxylate used as starting material in Example 5 is prepared by methods obvious to the chemist skilled in the art from the corresponding carboxylic acid prepared according to literature (Fuson et al., J. Org. Chem., 1948, 13, 489). Alkylating agents of formula VI are prepared from the corresponding dimethylamine (Formula VI:  $G = N(Me)_2$ ) as exemplified in example 9. The secondary amines of formula  $H-A-(CH_2)_m-B$  are commercially available, prepared by methods obvious to the chemist skilled in the art or prepared according to literature procedures. 1-(2-methoxyphenyl)piperazine is prepared according to Pollard et al., J. Org. Chem., 1958, 23, 1333. [2-(2-Methoxyphenoxy)ethyl]methylamine and [2-(3-methoxyphenoxy)ethyl]-methylamine are prepared as exemplified in Examples 7 and 10 using commercially available 2-methoxyphenoxyacetic acid and 3-methoxyphenoxyacetic acid, respectively, as starting materials.

15 The reductive alkylations according to method c and d are performed according to standard literature methods using  $NaCNBH_3$ ,  $NaBH_4$  or  $NaBH(OAc)_3$  as reducing agent in a suitable solvent.

The reductions according to Methods e and f are generally performed by use of  $LiAlH_4$ ,  $AlH_3$  or diborane in an inert solvent such as tetrahydrofuran, dioxane, or diethyl ether at room temperature or at a slightly elevated temperature.


The release of final products by means of Hofmann elimination in Method g is generally performed by the use of an organic base such as triethylamine or diisopropylethylamine in an aprotic organic solvent such as dichloromethane, toluene or N,N-dimethylformamide. The polymer of formula XII is prepared in a synthesis sequence as exemplified in Example 4 and described in the following. The starting acryl ester resin ( $CH_2CHC(O)OR'$ ) is prepared according to literature procedures (Brown et al., J. Am. Chem. Soc., 1997, 119, 3288-95) by acylation of commercially available hydroxy substituted resins such as cross linked hydroxymethylpolystyrene or Wang resin with acryloyl chloride. Secondary amines of formula  $H_2N-D-Z-(CH_2)_m-B$  are introduced by Michael addition in an organic solvent



such as N,N-dimethylformamide at ambient temperature. The secondary amines used are either commercially available, prepared by methods obvious to the chemist skilled in the art or prepared according to literature procedures. 3-(2-Methoxyphenyl)propylamine is prepared according to Leeson et al., J. Med. Chem. 1988, 31, 37-54, 3-(3-methoxyphenyl)propylamine according to Meise et al. Liebigs Ann. Chem., 1987, 639-42, 3-(2-methoxyphenoxy)propylamine according to Augsein et al., J. Med. Chem., 1965, 8, 356-67, 3-(3-methoxyphenoxy)propylamine according to Bremner et al., Aust. J. Chem. 1984, 37, 129-41, 2-benzyloxyethylamine according to Harder et al. Chem. Ber. 1964, 97, 510-19, 2-(1*H*-indolyl-3-yl)ethylamine according to Nenitzescu et al., Chem. Ber., 1958, 91, 1141-45 and 3-(1*H*-indolyl-3-yl)propylamine according to Jackson et al., J. Am. Chem. Soc., 1930, 52, 5029. The second diversifying group is introduced by means of alkylation with an agent of formula VI by boiling the reactants under reflux or by heating them at a fixed temperature in a suitable solvent such as tetrahydrofuran, dioxane, ethanol, 2-propanol, ethyl acetate, N,N-dimethylformamide, dimethyl sulfoxide or 1-methyl-2-pyrrolidinone in the presence of a soluble base such as diisopropylethylamine or triethylamine, or by means of reductive alkylation with an aldehyde of formula IX using standard solid phase synthesis literature methods using NaCNBH<sub>3</sub>, NaBH<sub>4</sub> or NaBH(OAc), as reducing agent in a suitable solvent. The third diversifying group was introduced by means of quarternisation using an alkylating agent of formula R<sup>4</sup>-G in an organic solvent such as tetrahydrofuran, dioxane, ethanol, 2-propanol, ethyl acetate, N,N-dimethylformamide, dimethyl sulfoxide or 1-methyl-2-pyrrolidinone at ambient temperature giving resins of formula XII.

The indole formation according to method h is performed by the reaction of acetals of formula XIII with aryl hydrazines of formula XIV resulting in the corresponding hydrazones, which subsequently are converted into indoles by means of the Fischer indole synthesis. The synthesis sequence is preferably performed as a one-pot procedure using a Lewis acid catalysts, preferably zinc chloride or boron fluoride, or protic acids, preferably sulfuric acid or phosphoric acid, in a suitable solvent such as acetic acid or ethanol at an elevated temperature. Acetals of formula XIII are prepared by alkylation of secondary amines of formula V with acetals of formula XV


16



XV

using the conditions described above for methods a and b. Alternatively, the acetals of formula XIII are prepared by alkylation of acetals of formula XVI

5



XVI

with an alkylating agent of formula VI using the conditions described above for methods a and b. The acetals of formula XVI are prepared by reaction of acetals of formula XV with primary amines of formula  $\text{NH}_2\text{R}^4$  using standard conditions.

Polymer bound acetals of formula XV is prepared by reaction of aldehydes of formula  $\text{G}-\text{B}-\text{CH}_2\text{CHO}$  with commercially available 2,2-dimethyl-1,3-dioxolan-4-yl-methoxymethyl polystyrene in a suitable solvent such as toluene, using p-toluenesulfonic acid as catalyst at elevated temperature. 4-Chlorobutanal, 5-chloropentanal, and 6-chlorohexanal were prepared in analogy to the method described by Normant et al., Tetrahedron 1994, 50 (40), 11665.

15

### Examples

Melting points were determined on a Büchi SMP-20 apparatus and are uncorrected. Mass spectra were obtained on a Quattro MS-MS system from VG Biotech, Fisons Instruments. The MS-MS system was connected to an HP 1050 modular HPLC system. A volume of 20-50  $\mu\text{l}$  of the sample (10  $\mu\text{g/ml}$ ) dissolved in a mixture of 1% acetic acid in acetonitril/water 1:1 was introduced via the autosampler at a flow of 30  $\mu\text{l/min}$  into the Electrospray Source. Spectra were obtained at two standard sets of operating conditions. Analytical LC-MS data were obtained on a PE Sciex API 150EX instrument equipped with IonSpray source and Shimadzu LC-8A/SLC-10A LC system. The LC conditions (50 X 4.6

25

mm YMC ODS-A with 5  $\mu$ m particle size) were linear gradient elution with water/acetonitrile/trifluoroacetic acid (90:10:0.05) to water/acetonitrile/trifluoroacetic acid (10:90:0.03) in 7 min at 2 mL/min. Purity was determined by integration of the UV trace (254 nm). The retention times  $R_t$  are expressed in minutes.

- 5 One set to obtain molecular weight information ( $MH^+$ ) (21 eV) and the other set to induce fragmentation patterns (70 eV). The background was subtracted. The relative intensities of the ions are obtained from the fragmentation pattern. When no intensity is indicated for the Molecular Ion ( $MH^+$ ), this ion was only present under the first set of operating conditions. Preparative LC-MS-separation was performed on the same instrument. The LC conditions
- 10 (50 X 20 mm YMC ODS-A with 5  $\mu$ m particle size) were linear gradient elution with water/acetonitrile/trifluoroacetic acid (80:20:0.05) to water/acetonitrile/trifluoroacetic acid (10:90:0.03) in 7 min at 22.7 mL/min. Fraction collection was performed by split-flow MS detection.

- $^1H$  NMR spectra were recorded at 500.13 MHz on a Bruker Avance DRX500 instrument or
- 15 at 250.13 MHz on a Bruker AC 250 instrument. Deuterated chloroform (99.8%D) or dimethyl sulfoxide (99.9%D) were used as solvents. TMS was used as internal reference standard. Chemical shift values are expressed in ppm-values. The following abbreviations are used for multiplicity of NMR signals: s=singlet, d=doublet, t=triplet, q=quartet, qui=quintet, h=heptet, dd=double doublet, dt=double triplet, dq=double quartet, tt=triplet
- 20 of triplets, m=multiplet. NMR signals corresponding to acidic protons are generally omitted. Content of water in crystalline compounds was determined by Karl Fischer titration. Standard workup procedures refer to extraction with the indicated organic solvent from proper aqueous solutions, drying of combined organic extracts (anhydrous  $MgSO_4$  or  $Na_2SO_4$ ), filtering and evaporation of the solvent *in vacuo*. For column chromatography
- 25 silica gel of type Kieselgel 60, 230-400 mesh ASTM was used.

### Example 1

- (+)-1-[3-[[4-(1,4-Benzodioxan-5-yl)butyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (1a). A mixture of 5-(4-bromobutyl)-1,4-
- 30 benzodioxane (1.5 g, 5.5 mmol), (+)-1-[3-(methylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (2.2. g, 5.5 mmol), potassium carbonate (3.0 g, 22

mmol), and methyl isobutyl ketone (150 mL) was boiled under reflux for 16 h. After cooling to room temperature the organic phase was washed with water (150 mL), the solvents evaporated *in vacuo* and the remaining oil purified by column chromatography (ethyl acetate/heptane/triethylamine 75:20:5) affording 2.0 g (73%) of the title compound as an oil:  $[\alpha]_D^{22} + 8.93^\circ$  (c 0.5; CH<sub>3</sub>OH). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.25-1.35 (m, 1H), 1.40-1.60 (m, 5H), 2.05-2.30 (m, 9H), 2.55 (t, 2H), 4.20-4.30 (m, 4H), 5.10-5.20 (m, 2H), 6.65-6.75 (m, 3H), 7.00 (t, 2H), 7.35 (d, 1H), 7.40 (dd, 2H), 7.50 (s, 1H), 7.60 (d, 1H); MS m/z 501 (MH<sup>+</sup>, 100), 262 (27), 149 (77), 109 (52).

10 The following compounds were prepared analogously:

(+)-1-[3-[[3-(1,4-Benzodioxan-5-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile oxalate (1b): mp 114-16°C (ethyl acetate);  $[\alpha]_D^{22} + 8.96^\circ$  (c 1.0; CH<sub>3</sub>OH); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$  1.35-1.45 (m, 1H), 1.45-1.55 (m, 1H), 1.80 (m, 2H), 2.20-2.30 (m, 2H), 2.45-2.55 (m, 2H), 2.60 (s, 3H), 2.90 (m, 2H), 2.95 (m, 2H), 4.20-4.30 (m, 4H), 5.20 (m, 2H), 6.65-6.75 (m, 3H), 7.10-7.20 (m, 2H), 7.55-7.60 (m, 2H), 7.70-7.80 (m, 1H), 7.80-7.95 (m, 2H); MS m/z 488 (MH<sup>+</sup>, 100), 262 (33), 149 (52), 109 (55).

1-[3-[[2-(1,4-Benzodioxan-5-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile oxalate (1c): mp 118-20°C (ethyl acetate); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$  1.40-1.70 (m, 2H), 2.25 (t, 2H), 2.70 (s, 3H), 2.75-2.90 (m, 2H), 2.90-3.15 (m, 4H), 4.15-4.30 (m, 4H), 5.20 (m, 2H), 6.65-6.80 (m, 3H), 7.20 (t, 2H), 7.60 (dd, 2H), 7.70-7.85 (m, 3H); MS m/z 473 (MH<sup>+</sup>, 64), 323 (13), 262 (24), 163 (100), 109 (25).

25 1-[3-[[1,4-Benzodioxan-5-yl)methyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile oxalate (1d): mp 160-62 °C (acetone/methanol); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$  1.40-1.70 (m, 2H), 2.25 (t, 2H), 2.60 (s, 3H), 2.90 (t, 2H), 4.00 (s, 2H), 4.20-4.30 (m, 4H), 5.20 (m, 2H), 6.80-7.00 (m, 3H), 7.15 (t, 2H), 7.50-7.65 (dd, 2H), 7.70-7.85 (m, 3H); MS m/z 459 (MH<sup>+</sup>, 7), 109 (100).

30

**Example 2**

1-(4-Fluorophenyl)-1-[3-[4-(2-methoxyphenyl)piperazinyl]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**2a**). A mixture of 1-(3-chloropropyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (2.5 g, 7.9 mmol), 1-(2-methoxyphenyl)piperazine (2.0 g, 10.4 mmol), potassium carbonate (3 g, 22 mmol) and methyl isobutyl ketone (200 mL) was boiled under reflux for 16 h. After cooling to room temperature the organic phase was washed with water (200 mL), the solvents were evaporated *in vacuo* and the remaining oil purified by column chromatography (ethyl acetate/heptane/triethylamine 75:20:5). The title compound crystallised from diethyl ether

1.5 g (40 %): mp 147-49 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 1.30-1.65 (m, 2H), 2.10-2.30 (m, 2H), 2.40 (t, 2H), 2.50-2.70 (m, 4H), 2.90-3.20 (m, 4H), 3.85 (s, 3H), 5.20 (m, 2H), 6.70-7.10 (m, 6H), 7.30-7.55 (m, 4H), 7.60 (d, 1H); MS *m/z*, 472 (MH<sup>+</sup>, 100), 262 (14), 109 (19).

The following compounds were prepared analogously:

1-(4-Fluorophenyl)-1-[3-[[2-(2-methoxyphenoxy)ethyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**2b**): (oil) <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.30-1.40 (m, 1H), 1.40-1.55 (m, 1H), 2.10-2.20 (m, 2H), 2.25 (s, 3H), 2.40-2.45 (t, 2H), 2.70-2.80 (m, 2H), 3.70 (s, 3H), 4.05 (t, 2H), 5.15 (m, 2H), 6.85-7.00 (m, 6H), 7.30-7.45 (m, 3H), 7.50 (s, 1H), 7.55 (d, 1H).

1-(4-Fluorophenyl)-1-[3-[[2-(3-methoxyphenoxy)ethyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**2c**): (oil) <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.30-1.40 (m, 1H), 1.40-1.55 (m, 1H), 2.10-2.20 (m, 2H), 2.25 (s, 3H), 2.40 (t, 2H), 2.70-2.75 (m, 2H), 3.70 (s, 3H), 4.00 (t, 2H), 5.15 (m, 2H), 6.40-6.55 (m, 3H), 7.00 (t, 2H), 7.20 (t, 1H), 7.35 (d, 1H), 7.40 (dd, 2H), 7.50 (s, 1H), 7.55 (d, 1H).

(*S*)-1-[3-[[4-(1*H*-Indol-3-yl)butyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**2d**): LC/MS (*m/z*) 482 (MH<sup>+</sup>), *R*<sub>t</sub> = 4.24, purity: 84 %.

1-[3-[[4-(1*H*-Indol-3-yl)butyl]methylamino]propyl]-1-phenyl-1,3-dihydroisobenzofuran (**2e**): LC/MS (*m/z*) 439 (MH<sup>+</sup>), *R*<sub>t</sub> = 4.33, purity: 77 %.

- (S)-1-[3-[[3-(1*H*-Indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**2f**): LC/MS (*m/z*) 468 (*MH*<sup>+</sup>), *Rt* = 4.11, purity: >99 %.
- 1-3-[[3-(1*H*-Indol-3-yl)propyl]methylamino]propyl]-1-phenyl-1,3-dihydroisobenzofuran (**2g**): LC/MS (*m/z*) 425 (*MH*<sup>+</sup>), *Rt* = 4.15, purity: >99 %.
- 5-3-[[3-(1-Phenyl-1,3-dihydroisobenzofuran-1-yl)propyl]methylamino]propyl]-1,4-benzodioxane (**2h**): LC/MS (*m/z*) 444 (*MH*<sup>+</sup>), *Rt* = 4.12, purity: 97 %.
- 5-3-[[3-[1-(3-Chlorophenyl)-1,3-dihydroisobenzofuran-1-yl]propyl]methylamino]propyl]-1,4-benzodioxane (**2i**): LC/MS (*m/z*) 478 (*MH*<sup>+</sup>), *Rt* = 4.45, purity: 93 %.
- 10 5-3-[[3-[1-(4-Fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]propyl]methylamino]propyl]-1,4-benzodioxane (**2j**): LC/MS (*m/z*) 462 (*MH*<sup>+</sup>), *Rt* = 4.21, purity: 93 %.
- 5-3-[[3-[1-(3-Trifluoromethylphenyl)-1,3-dihydroisobenzofuran-1-yl]propyl]methylamino]propyl]-1,4-benzodioxane (**2k**): LC/MS (*m/z*) 512 (*MH*<sup>+</sup>), *Rt* = 4.59, purity: 90 %.
- 15 1-3-[[3-(1,4-Benzodioxan-5-yl)propyl]methylamino]propyl]-1-(4-chlorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**2l**): LC/MS (*m/z*) 503 (*MH*<sup>+</sup>), *Rt* = 4.59, purity: >99 %.
- 1-3-[4-(1*H*-Indol-4-yl)piperazinyl]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**2m**): LC/MS (*m/z*) 481 (*MH*<sup>+</sup>), *Rt* = 5.61, purity: 97 %.
- 20 1-3-[4-(1*H*-Indol-5-yl)piperazinyl]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**2n**): LC/MS (*m/z*) 481 (*MH*<sup>+</sup>), *Rt* = 5.69, purity: 94 %.
- 1-3-[4-(6-chloro-1*H*-Indol-3-yl)piperidinyl]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**2o**): LC/MS (*m/z*) 514 (*MH*<sup>+</sup>), *Rt* = 6.38, purity: 96 %.

25

**Example 3**

- 5-3-[[3-[-5-Fluoro-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]propyl]methylamino]propyl]-1,4-benzodioxane oxalate (**3**). A solution of 3-(1,4-benzodioxan-5-yl)propionic acid (0.8 g, 3.8 mmol), thionyl chloride (1 mL, 13.7 mmol)
- 30 and one droplet of *N,N*-dimethylformamide in dichloromethane (30 mL) was boiled under reflux for 2 h. The volatile solvents were evaporated *in vacuo* and the remaining oil was

dissolved in dichloromethane (30 mL). The resulting solution was added to a solution [3-  
5-fluoro-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]propyl]methylamine (3.0 g, 10  
mmol) and triethylamine (10 mL) in dichloromethane (100 mL). After stirring for 16 h the  
volatile solvents were evaporated *in vacuo* and the remaining oil was purified by column  
5 chromatography (ethyl acetate/heptane 75:25) affording 1.4 g of crude amide which was  
used without further purification.

To a solution of the amide (1.4 g, 2.8 mmol) in tetrahydrofuran (200 mL) was added  
lithium aluminum hydride (1.0 g, 2.6 mmol). After boiling of the resulting mixture under  
reflux for 3 h, the reaction mixture was cooled to 0 °C and carefully treated with water (1  
10 mL) and 4 N aqueous sodium hydroxide (1 mL). The resulting mixture was filtered and  
dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the volatile solvents afforded the title compound as an oil  
which was precipitated as its oxalate in acetone 0.9 g (19%): mp 131-33 °C; <sup>1</sup>H NMR  
(DMSO-*d*<sub>6</sub>) δ 1.35-1.45 (m, 1H), 1.45-1.55 (m, 1H), 1.75-1.80 (m, 2H), 2.10-2.25 (m, 2H),  
2.50-2.55 (m, 2H), 2.60 (s, 3H), 2.90 (t, 2H), 2.95 (t, 2H), 4.20-4.25 (m, 4H), 5.10 (m,  
15 2H), 6.65-6.75 (m, 3H), 7.10-7.15 (m, 4H), 7.45-7.60 (m, 3H); MS *m/z*, 480 (MH<sup>+</sup>, 100),  
225 (34), 109 (51).

#### Example 4

1-[3-[[2-(1*H*-Indol-3-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-  
20 dihydroisobenzofuran-5-carbonitrile (4a). To a suspension of acryl ester Wang resin  
(CH<sub>2</sub>CHC(O)OR', HOR' = Wang resin) (loading 1.0 mmol/g) (300 mg, 0.30 mmol)  
(prepared from Wang resin (Loading 1.09 mmol/g, 200-400 mesh, 1% divinylbenzene) in  
analogy with the procedure described for the preparation of acryl ester hydroxymethyl  
polystyrene by Brown et al., J. Am. Chem. Soc., 1997, 119, 3288-95) in N,N-  
25 dimethylformamide (1.5 mL) was added a solution of 2-(1*H*-indolyl-3-yl)ethylamine (96  
mg, 0.60 mmol) in N,N-dimethylformamide (1.5 mL). After stirring of the resulting  
suspension at room temperature for 16 h, the resin was filtered off and subsequently  
washed with 0.3 M diisopropylethylamine in N,N-dimethylformamide (3 X 2.5 mL),  
methanol (2 X 2.5 mL) and dichloromethane (2 X 2.5 mL).  
30 To a suspension of the resulting resin in acetonitrile (1.5 mL) was added a solution of 1-(3-  
chloropropyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (9)

(473 mg, 1.5 mmol) in acetonitrile (1.5 mL) and diisopropylethylamine (280 mL, 1.6 mmol). After heating of the resulting mixture at 75 °C under stirring for 16 h, the resin was filtered off. The resin was subsequently washed with acetonitrile (3 X 2.5 mL), methanol (3 X 2.5 mL), and dichloromethane (3 X 2.5 mL). The resin was suspended in N,N-dimethylformamide and diisopropylethylamine (280 mL, 1.6 mmol) and acetic anhydride (140 mL, 1.5 mmol) was added. After stirring of the resulting mixture for 16 h the resin was filtered off and washed with N,N-dimethylformamide (3 X 2.5 mL), methanol (3 X 2.5 mL), and dichloromethane (3 X 2.5 mL).

The intermediate resin was suspended in N,N-dimethylformamide (2 mL) and a solution of iodomethan (187 mL, 3.0 mmol) in N,N-dimethylformamide was added. After stirring of the resulting mixture for 16 h at room temperature, the resin was filtered off and washed with N,N-dimethylformamide (3 X 2.5 mL), methanol (3 X 2.5 mL), and dichloromethane (3 X 2.5 mL). To the resulting resin was added N,N-dimethylformamide (3.0 mL) and Diisopropylethylamine (165 mL, 0.94 mmol) and the mixture was stirred for 16 h. The resin was filtered off and washed with methanol (2 X 2.0 mL). The cleavage solution and the washing solutions were collected and the solvent evaporated *in vacuo*. The remaining oil was purified by ion exchange chromatography using an 6 mL Varian SCX column (1225-6011). The column was preconditioned with 10% acetic acid in methanol (3 mL) and the crude product was loaded on the column in a 2:1 mixture of methanol and 1-methyl-2-pyrrolidinone (3 mL). After the column was washed with methanol (18 mL) and acetonitrile (3 mL) the product was eluted from the column with 4 N ammonia in methanol (4 mL) and subsequent evaporation of the solvents *in vacuo* afforded 13.9 mg (10%) of the title compound as an oil: LC/MS (m/z) 454 (MH<sup>+</sup>), Rt = 6.13 , purity: 98 %.

The following compounds were prepared analogously:

1-(4-Fluorophenyl)-1-[3-[[2-(3-methoxyphenyl)ethyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**4b**): LC/MS (m/z) 445 (MH<sup>+</sup>), R<sub>t</sub> = 8.58 , purity: 88%

1-(4-Fluorophenyl)-1-[3-[[2-(3-methoxyphenyl)ethyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**4c**): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.30-1.60 (m, 2H), 2.00-2.20 (m, 2H), 2.40-2.55 (m, 2H), 2.55-2.70 (m, 4H), 3.00-3.15 (broad s, 2 H), 3.80 (s, 3H), 5.05-



5.20 (m, 4H), 5.75-5.85 (m, 1H), 6.65-6.80 (m, 3H), 7.00 (t, 2H), 7.20 (t, 1H), 7.30 (d, 1H), 7.40 (m, 2H), 7.50 (s, 1H), 7.60 (d, 1H); LC/MS (m/z) 471 (MH<sup>+</sup>), R<sub>t</sub> = 8.85 , purity: 91%

1-(4-Fluorophenyl)-1-[3-[[2-(2-methoxyphenyl)ethyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (4d): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.25-1.40 (m, 1H), 1.40-1.55 (m, 1H), 2.05-2.25 (m, 2H), 2.40-2.50 (m, 2H), 2.50-2.65 (m, 2H), 2.65-2.75 (m, 2H), 3.00-3.15 (m, 2 H), 3.80 (s, 3H); 5.05-5.20 (m, 4H), 5.75-5.90 (m, 1H), 6.75-6.90 (m, 2H), 6.95-7.10 (m, 3H), 7.20 (t, 1H), 7.30 (d, 1H), 7.35-7.45 (m, 2H), 7.45 (s, 1H), 7.60 (d, 1H); LC/MS (m/z) 471 (MH<sup>+</sup>), R<sub>t</sub> = 7.82 , purity: >89%

10

1-[3-[[2-(2,5-Dimethoxyphenyl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (4e): LC/MS (m/z) 475 (MH<sup>+</sup>), R<sub>t</sub> = 8.68 , purity: 94%

1-[3-[[2-(2,5-Dimethoxyphenyl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (4f): LC/MS (m/z) 500 (MH<sup>+</sup>), R<sub>t</sub> = 8.95 , purity: 90%

1-(4-Fluorophenyl)-1-[3-[[2-phenoxyethyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (4g): LC/MS (m/z) 431 (MH<sup>+</sup>), R<sub>t</sub> = 8.58 , purity: 95%

20

1-[3-[[2-(1*H*-Indolyl-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (4h): LC/MS (m/z) 480 (MH<sup>+</sup>), R<sub>t</sub> = 8.87 , purity: 93%

1-(4-Fluorophenyl)-1-[3-[[2-phenoxyethyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (4i): LC/MS (m/z) 457 (MH<sup>+</sup>), R<sub>t</sub> = 6.40 , purity: >99%

25

1-(4-Fluorophenyl)-1-[3-[[3-(2-methoxyphenyl)propyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (4j): LC/MS (m/z) 459 (MH<sup>+</sup>), R<sub>t</sub> = 6.43 , purity: >99%

30

1-(4-Fluorophenyl)-1-[3-[[3-(2-methoxyphenyl)propyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**4k**): LC/MS (m/z) 485 (MH<sup>+</sup>), R<sub>t</sub> = 6.77 , purity: >99%

5 1-(4-Fluorophenyl)-1-[3-[[3-(3-methoxyphenyl)propyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**4l**): LC/MS (m/z) 485 (MH<sup>+</sup>), R<sub>t</sub> = 6.63 , purity: >99%

1-(4-Fluorophenyl)-1-[3-[[3-(2-methoxyphenoxy)propyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**4m**): LC/MS (m/z) 475 (MH<sup>+</sup>), R<sub>t</sub> = 6.20 , purity: >99%

1-(4-Fluorophenyl)-1-[3-[[3-(2-methoxyphenoxy)propyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**4n**): LC/MS (m/z) 501 (MH<sup>+</sup>), R<sub>t</sub> = 6.50 , purity: >99%

1-(4-Fluorophenyl)-1-[3-[[3-(3-methoxyphenoxy)propyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**4o**): LC/MS (m/z) 475 (MH<sup>+</sup>), R<sub>t</sub> = 6.35 , purity: >99%

20 1-(4-Fluorophenyl)-1-[3-[[3-(3-methoxyphenoxy)propyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile (**4p**): LC/MS (m/z) 501 (MH<sup>+</sup>), R<sub>t</sub> = 6.65 , purity: >99%

25 1-[3-[(2-Benzyloxyethyl)methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**4q**): LC/MS (m/z) 445 (MH<sup>+</sup>), R<sub>t</sub> = 6.18 , purity: 98%

1-[3-[(2-Benzyloxyethyl)(prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**4r**): LC/MS (m/z) 471 (MH<sup>+</sup>), R<sub>t</sub> = 6.55 , purity: 97%

30

1-[3-[[3-(1*H*-Indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**4s**): LC/MS (*m/z*) 468 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 6.28, purity: 80%

1-[3-[[3-(1*H*-Indol-3-yl)propyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**4t**): LC/MS (*m/z*) 494 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 6.60, purity: 82%

1-[3-[[3-(1*H*-Indol-3-yl)propyl](prop-2-yn-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**4u**): LC/MS (*m/z*) 492 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 6.59, purity: 73%

#### 10 Example 5

5-hydroxymethyl-1,4-benzodioxan (**5**). To a suspension of lithium aluminum hydride (7.0 g, 0.18 mol) in dry diethyl ether (100 mL) was added a solution of ethyl 1,4-benzodioxan-5-carboxylate (35 g, 0.17 mol) in diethyl ether (100 mL). After boiling under reflux for 2 h, the reaction mixture was cooled to 0 °C and carefully treated with water (35 mL) and 4 N aqueous sodium hydroxide (35 mL). The resulting mixture was filtered and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvents afforded 25 g (88%) crystalline title compound: mp 51-53 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 2.50 (s, 1H), 4.20-4.3 (m, 4H), 4.60 (s, 2H), 6.75-6.90 (m, 3H).

#### Example 6

20 2-(1,4-benzodioxan-5-yl)acetic acid (**6**). To a solution of 5-hydroxymethyl-1,4-benzodioxan (8.0 g, 48 mmol) in dichloromethane (200 mL) was added two droplets of N,N-dimethylformamide and thionyl chloride (5.0 mL, 68 mmol) at room temperature. After the resulting solution was boiled under reflux for 1 h and subsequently cooled to room temperature water (100 mL) was added. The phases were separated and the organic phase was dried (MgSO<sub>4</sub>) and the solvents evaporated *in vacuo*. A solution of the remaining oil (8.5 g, 46 mmol) was added to a mixture of sodium cyanide (5.0 g, 102 mmol) and N,N-dimethylformamide (100 mL) at room temperature. After stirring for 16 h at room temperature ice was added and the resulting slurry was extracted with diethyl ether (2 X 250 mL). The collected organic phases were washed with saturated calcium chloride, dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvents were evaporated *in vacuo*. A mixture of the remaining oil 30 (6.0 g, 34 mmol), ethanol (200 mL), sodium hydroxide (6.0 g) and water (6 mL) was

boiled under reflux for 16 h. After evaporation of the solvents *in vacuo*, water (200 mL) was added and the resulting slurry was extracted with diethyl ether (2 X 200 mL). The collected organic phases were washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ) and the solvents were evaporated *in vacuo* affording 4.0 g (43%) of the title compound as an oil:  $^1\text{H}$  NMR

5 (CDCl<sub>3</sub>)  $\delta$  3.65 (s, 2H), 4.15-4.30 (m, 4H), 6.70-6.85 (m, 3H).

#### Example 7

5-(2-bromoethyl)-1,4-benzodioxan (7a). To a solution of 2-(1,4-benzodioxan-5-yl)acetic acid (6) (4.0 g, 21 mmol) in tetrahydrofuran (200 mL) was added lithium aluminum hydride (1.0 g, 26 mmol). After boiling under reflux for 2 h the reaction mixture was cooled to 0 °C and carefully treated with water (1 mL) and 4 N aqueous sodium hydroxide (1 mL). The resulting mixture was filtered and dried ( $\text{Na}_2\text{SO}_4$ ). Evaporation of the solvents afforded crude intermediate alcohol (3.9 g, 21 mmol) as an oil which was used without further purification. To a solution of the intermediate alcohol and tetrabromomethane (8.8 g, 27 mmol) in acetonitrile (120 mL) was added triphenylphosphine (6.3 g, 24.9 mmol) in small portions at 0 °C. After reaction for further 15 minutes at 0 °C the solvents were evaporated *in vacuo* and the remaining oil was purified by column chromatography (ethyl acetate/heptane 66:34) affording 5.5 g (99%) of the title compound as an oil:  $^1\text{H}$  NMR (CDCl<sub>3</sub>)  $\delta$  3.15 (t, 2H), 3.55 (t, 2H), 4.20-4.35 (m, 4H), 6.65-6.85 (m, 3H).

The following compounds were prepared analogously:

5-(3-Bromopropyl)-1,4-benzodioxan (7b): (oil)  $^1\text{H}$  NMR (CDCl<sub>3</sub>)  $\delta$  2.15 (qui, 2H), 2.75 (t, 2H), 3.40 (t, 2H), 4.20-4.30 (m, 4H), 6.65-6.75 (m, 3H).

5-(4-Bromobutyl)-1,4-benzodioxan (7c): (oil)  $^1\text{H}$  NMR (CDCl<sub>3</sub>)  $\delta$  1.70-1.80 (qui, 2H), 1.85-1.90 (qui, 2H), 2.60 (t, 2H), 3.40 (t, 2H), 4.25 (m, 4H), 6.65-6.75 (m, 3H).

1-(2-Bromoethoxy)-2-methoxybenzene (7d): (oil)  $^1\text{H}$  NMR (CDCl<sub>3</sub>)  $\delta$  3.65 (t, 2H), 3.85 (s, 3H), 4.30 (t, 2H), 6.80-7.05 (m, 4H).

1-(2-Bromoethoxy)-3-methoxybenzene (7e): (oil) <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.60 (t, 2H), 3.80 (s, 3H), 4.25 (t, 2H), 6.45-6.55 (m, 3H), 7.15 (t, 1H).

### Example 8

- 5 4-(1,4-Benzodioxan-5-yl)butanoic acid (8a). Neat 5-(4-bromoethyl)-1,4-benzodioxan (7c) (18.0 g, 74 mmol) was added to a mixture of diethyl malonate (12 g, 75 mmol), potassium tert-butoxide (8.4 g, 75 mmol), toluene (250 mL) and dimethyl sulfoxide (50 mL) at room temperature. The resulting mixture was heated at 50 °C for 3 h, cooled to room temperature and water was added. After the slurry was acidified with concentrated hydrochloric acid the
- 10 phases were separated. The organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvents evaporated *in vacuo*. The remaining oil was dissolved in ethanol (200 mL) and 9 N aqueous sodium hydroxide. After boiling of the resulting mixture under reflux for 15 minutes the solution was stirred at room temperature for 1 h. The solvents were evaporated and the remaining oil was diluted in water (200 mL) and extracted with diethyl ether (2 X 100 mL). The
- 15 aqueous phase was acidified with 4 N hydrochloric acid and extracted with ethyl acetate (2 X 200 mL). Drying of the collected organic phases and evaporation of the solvents *in vacuo* afforded the intermediate dicarboxylic acid as an oil (5.0 g). The crude oil was diluted in pyridine (10 mL) and the resulting solution was heated at 115 °C for 1 h. After cooling to room temperature, water (50 mL) was added and the aqueous phase was
- 20 acidified with 4 N hydrochloric acid. The resulting slurry was extracted with diethyl ether (2 X 50 mL) and the collected organic phases were dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvents *in vacuo* afforded 3.8 g (23%) of the title compound as an oil.

The following compound was prepared analogously:

- 25 3-(1,4-Benzodioxan-5-yl)propionic acid (8b): (oil) <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 2.65 (t, 2H), 2.95 (t, 2H), 4.20-4.30 (m, 4H), 6.65-6.80 (m, 3H).

### Example 9

- 1-(3-Chloropropyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (9). To a
- 30 mixture of 1-[3-(methylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (43 g, 138 mmol), potassium carbonate (30 g, 217 mmol), and ethanol (400

mL) was added ethyl bromoacetate (20 mL, 180 mmol) at room temperature and the resulting mixture was boiled under reflux for 90 minutes. After cooling to room temperature water (800 mL) and ethyl acetate (500 mL) was added and the phases were separated. The organic phase was washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ) and the solvents  
5 evaporated *in vacuo*. The remaining oil (36 g, 101 mmol) was added slowly to a mixture of ethyl chloroformate (50 mL, 523 mmol), potassium carbonate (36 g, 260) and toluene (300 mL) at 90 °C. After boiling of the resulting mixture under reflux for 1 h and cooling to room temperature, the solvents were evaporated *in vacuo*. The remaining oil was purified by column chromatography (ethyl acetate/heptane 1:3) giving 15 g (34%) of the title  
10 compound as an oil:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.60-1.90 (m, 2H), 2.20-2.45 (m, 2H), 3.45-3.55 (m, 2H), 5.20 (m, 2H), 6.95-7.10 (t, 2H), 7.40-7.55 (m, 4H), 7.60 (d, 1H).

#### Example 10

[2-(2-Methoxyphenoxy)ethyl]methylamine (10a). A solution of 1-(2-bromoethoxy)-2-  
15 methoxybenzene (7d) (7.7 g, 33 mmol) in a 33% solution of methylamine in ethanol was heated at 80 °C in a sealed tube for 16 h. After cooling to room temperature, the solvents were evaporated *in vacuo*. A 2 N aqueous solution of sodium hydroxide was added to the remaining oil and the resulting slurry was extracted with ethyl acetate (2 X 250 mL). The collected organic phases were dried ( $\text{Na}_2\text{SO}_4$ ) and the solvents evaporated *in vacuo* giving  
20 5.9 g (98%) of the title compound as an oil:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.85 (broad s, 1H), 2.50 (s, 3H), 3.00 (t, 2H), 3.85 (s, 3H), 4.10 (t, 2H), 6.85-6.95 (m, 4H).

The following compound was prepared analogously:

[2-(3-Methoxyphenoxy)ethyl]methylamine (10b): (oil)  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.85 (broad s,  
25 1H), 2.50 (s, 3H), 2.95 (t, 2H), 3.80 (s, 3H), 4.05 (t, 2H), 6.45-6.55 (m, 3H), 7.15 (t, 1H).

#### Example 11

2-(4-Chlorobutyl)-dioxolan-4-ylmethoxymethyl polystyrene (11a). A 2 L round bottom flask was charged with 2,2-dimethyldioxolan-4-ylmethoxymethyl polystyrene (90 g, 72  
30 mmol, commercially available as ( $\pm$ )-1-(2,3-isopropylidene) glycerol polystyrene from Calbiochem-Novabiochem, cat. no. 01-64-0291). Toluene (900 mL) followed by p-

29.

toluenesulfonic acid mono hydrate (5.0 g, 26 mmol), sodium sulfate (25 g), and readily available 5-chloropentanal (25.5 g, 211 mmol) were added and the mixture heated at reflux for 12 h. The reflux condenser was replaced by a Dean-Stark apparatus and the mixture was heated at reflux for an additional 3 h. After cooling of the reaction mixture to 60 °C, the resin was filtered and washed with toluene (200 mL), tetrahydrofuran/pyridine (1:1, 200 mL), tetrahydrofuran/water/pyridine (10:10:1, 200 mL), methanol (200 mL), water (200 mL), tetrahydrofuran (200 mL), dichloromethane (200 mL), methanol (3 X 200 mL), and dichloromethane (3 X 200 mL). The resin was dried *in vacuo* (55 °C, 12 h) to yield the title compound 11a (97 g).

The following compounds were prepared analogously:

2-(3-Chloropropyl)-dioxolan-4-ylmethoxymethyl polystyrene (11b)

2-(5-Chloropentyl)-dioxolan-4-ylmethoxymethyl polystyrene (11c)

## Example 12

1-[3-[[3-(1*H*-Indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (4s). 2-(4-Chlorobutyl)-dioxolan-4-ylmethoxymethyl polystyrene (11a) (8.0 g, 6.1 mmol) was suspended in dry N,N-dimethylformamide (90 mL). Sodium iodide (3.38 g, 22.5 mmol) was added followed by diisopropylethylamine (6.30 mL, 36 mmol) and 1-[3-(methylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (5.56 g, 18 mmol). The reaction mixture was heated at 80 °C under stirring for 12 h. After cooling to room temperature, the resin was filtered and washed with N,N-dimethylformamide (3 X 65 mL), methanol (3 X 60 mL), tetrahydrofuran (3 X 60 mL), and then subsequently with methanol and tetrahydrofuran (each approximately 40 mL, 5 cycles). Finally, the resin was washed with tetrahydrofuran (4 X 40 mL) and dried *in vacuo* (55 °C, 12 h, 9.5 g).

An aliquot of this material (147 mg, 0.112 mmol) and phenylhydrazine hydrochloride (43 mg, 0.297 mmol) were mixed in a reactor tube. A 0.5 M solution of anhydrous zinc chloride in acetic acid (1.5 mL) was added and the reaction tube was sealed. The reaction mixture was stirred for 12 h at 75 °C. After cooling to room temperature, the reaction mixture was filtered and the residual resin washed with dimethylsulfoxide (1.5 mL). To the combined filtrates was added saturated aqueous sodium bicarbonate solution (1.5 mL). The

solution was loaded on a reversed solid phase extraction column (C-18, 1 g, Varian Mega Bond Elut®, Chrompack cat. no. 220508), pre-conditioned with methanol (3 mL) and water (3 mL). The column was washed with water (4 mL) and the product was eluted with methanol (4.5 mL). The resulting solution was loaded on an ion exchange column (SCX, 1 g, Varian Mega Bond Elut®, Chrompack cat. no. 220776), pre-conditioned with 10 % solution of acetic acid in methanol (3 mL) and the column was washed with methanol (4 mL) and acetonitrile (4 mL), followed by elution with 4 N solution of ammonia in methanol (4.5 mL). Evaporation of the volatile solvents afforded the title compound (4s) as a colourless oil (22 mg, 42 %). LC/MS (m/z) 468 (MH<sup>+</sup>), Rt = 4.30, purity: 83 %.

10

The following compounds were prepared analogously:

1-[3-[[2-(5-Methyl-1*H*-indol-3-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (12a): LC/MS (m/z) 468 (MH<sup>+</sup>), Rt = 4.22, purity: 96 %.

15

1-[3-[[2-(7-Fluoro-1*H*-indol-3-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (12b): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.2-1.4 (m, 1H), 1.4-1.55 (m, 1H), 2.0-2.25 (m, 2H), 2.25 (s, 3H), 2.39 (t, 2H), 2.60 (t, 2H), 2.86 (t, 2H), 5.05-5.21 (m, 2H), 6.93-7.07 (m, 4H), 7.17-7.3 (m, 2H), 7.3-7.4 (m, 3H), 7.4-7.5 (m, 1H), 7.5-7.6 (m, 1H); LC/MS (m/z) 472 (MH<sup>+</sup>), Rt = 4.12, purity: 86 %.

20

5-Fluoro-1-[3-[[3-(5-methyl-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran (12c): LC/MS (m/z) 475 (MH<sup>+</sup>), Rt = 4.57, purity: 92 %.

5-Fluoro-1-[3-[[3-(7-fluoro-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran (12d): LC/MS (m/z) 479 (MH<sup>+</sup>), Rt = 4.47, purity:

25

94 %.

1-[3-[[3-(5-Methyl-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (12e): LC/MS (m/z) 482 (MH<sup>+</sup>), Rt = 4.54, purity: 80 %.

1-[3-[Ethyl[3-(1*H*-indol-3-yl)propyl]amino]propyl]-1-(4-fluorophenyl)-1,3-

30

dihydroisobenzofuran-5-carbonitrile (12f): LC/MS (m/z) 482 (MH<sup>+</sup>), Rt = 4.31, purity: 94 %.



1-[3-[Ethyl[2-(5-methyl-1*H*-indol-3-yl)ethyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12g**): LC/MS (*m/z*) 482 (*MH*<sup>+</sup>), *Rt* = 4.38, purity: 89 %.

1-[3-[[3-(7-Fluoro-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12h**): LC/MS (*m/z*) 486 (*MH*<sup>+</sup>), *Rt* = 4.16, purity: 79 %.

1-[3-[[3-(5-Fluoro-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12i**): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.23-1.39 (m, 1H), 1.39-1.54 (m, 1H), 1.80 (tt, 2H), 2.06-2.24 (m, 5H), 2.30 (t, 2H), 2.34 (t, 2H), 2.68 (t, 2H), 5.13 (d, 1H), 5.17 (d, 1H), 6.93 (dt, 2H), 6.99 (t, 2H), 7.21 (dd, 1H), 7.23-7.29 (m, 1H), 7.33 (d, 1H), 7.40 (dd, 2H), 7.47 (s, 1H), 7.55 (d, 1H), 8.01 (s, 1H); LC/MS (*m/z*) 486 (*MH*<sup>+</sup>), *Rt* = 4.12, purity: 98 %.

1-[3-[Ethyl[2-(5-fluoro-1*H*-indol-3-yl)ethyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12j**): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.02 (t, 3H), 1.25-1.38 (m, 1H), 1.42-1.54 (m, 1H), 2.10 (ddd, 1H), 2.18 (ddd, 1H), 2.49 (t, 2H), 2.56 (q, 2H), 2.61-2.70 (m, 2H), 2.74-2.82 (m, 2H), 5.13 (d, 1H), 5.18 (d, 1H), 6.94 (dt, 2H), 6.99 (t, 2H), 7.19 (dd, 1H), 7.23-7.30 (m, 2H), 7.38 (dd, 2H), 7.47 (s, 1H), 7.54 (d, 1H), 8.01 (s, 1H); LC/MS (*m/z*) 486 (*MH*<sup>+</sup>), *Rt* = 4.24, purity: 95 %.

1-[3-[Ethyl[2-(7-fluoro-1*H*-indol-3-yl)ethyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12k**): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.02 (t, 3H), 1.22-1.37 (m, 1H), 1.42-1.53 (m, 1H), 2.0-2.2 (m, 2H), 2.36-2.6 (m, 4H), 2.67 (t, 2H), 2.81 (t, 2H), 5.12 (dd, 1H), 5.16 (d, 1H), 6.86-7.06 (m, 4H), 7.2-7.4 (m, 5H), 7.46 (d, 1H), 7.54 (d, 1H); LC/MS (*m/z*) 486 (*MH*<sup>+</sup>), *Rt* = 4.26, purity: 91 %.

1-[3-[[2-(5-Chloro-1*H*-indol-3-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12l**): LC/MS (*m/z*) 488 (*MH*<sup>+</sup>), *Rt* = 4.30, purity: 85 %.

1-[3-[[3-(5-Chloro-1*H*-indol-3-yl)propyl]methylamino]propyl]-5-fluoro-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran (**12m**): LC/MS (*m/z*) 495 (*MH*<sup>+</sup>), *Rt* = 4.64, purity: 94 %.

1-[3-[[4-(5-Methyl-1*H*-indol-3-yl)butyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12n**): LC/MS (*m/z*) 496 (*MH*<sup>+</sup>), *Rt* = 4.50, purity: 78 %.

1-[3-[Ethyl[3-(5-methyl-1*H*-indol-3-yl)propyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12o**): LC/MS (*m/z*) 496 (*MH*<sup>+</sup>), *Rt* = 4.50, purity: 92 %.

1-[3-[Ethyl[3-(7-fluoro-1*H*-indol-3-yl)propyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12p**): LC/MS (*m/z*) 500 (*MH*<sup>+</sup>), *Rt* = 4.39, purity: 91 %.

1-[3-[Ethyl[3-(5-fluoro-1*H*-indol-3-yl)propyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12q**): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.95 (t, 3H), 1.21-1.36 (m, 1H), 1.36-1.50 (m, 1H), 1.77 (tt, 2H), 2.10 (ddd, 1H), 2.18 (ddd, 1H), 2.34-2.50 (m, 6H), 2.65 (t, 2H), 5.12 (d, 1H), 5.15 (d, 1H), 6.90-7.04 (m, 4H), 7.20 (dd, 1H), 7.25 (dd, 1H), 7.30 (d, 1H), 7.36 (m, 2H), 7.45 (s, 1H), 7.52 (d, 1H), 8.12 (s, 1H); LC/MS (*m/z*) 500 (*MH*<sup>+</sup>), *Rt* = 4.35, purity: 94 %.

1-[3-[[3-(5-Chloro-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12r**): LC/MS (*m/z*) 502 (*MH*<sup>+</sup>), *Rt* = 4.55, purity: 91 %.

1-[3-[[2-(7-Chloro-1*H*-indol-3-yl)ethyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12s**): LC/MS (*m/z*) 502 (*MH*<sup>+</sup>), *Rt* = 4.41, purity: 80 %.

1-[3-[[2-(5-Chloro-1*H*-indol-3-yl)ethyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12t**): LC/MS (*m/z*) 502 (*MH*<sup>+</sup>), *Rt* = 4.44, purity: 95 %.

1-[3-[[2-(5,7-Difluoro-1*H*-indol-3-yl)ethyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12u**): LC/MS (*m/z*) 504 (*MH*<sup>+</sup>), *Rt* = 4.35, purity: 92 %.

- 1-[3-[[4-(5-Fluoro -1*H*-indol-3-yl)butyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12v**): LC/MS (*m/z*) 514 (*MH*<sup>+</sup>), *Rt* = 4.50, purity: 91 %.
- 5 1-[3-[[4-(5-Chloro -1*H*-indol-3-yl)butyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12w**): LC/MS (*m/z*) 516 (*MH*<sup>+</sup>), *Rt* = 4.59, purity: 90 %.
- 1-[3-[[3-(5-Chloro-1*H*-indol-3-yl)propyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12x**): LC/MS (*m/z*) 516 (*MH*<sup>+</sup>), *Rt* = 4.56, purity: 97 %.
- 10 1-[3-[[3-(5,7-Difluoro -1*H*-indol-3-yl)propyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12y**): LC/MS (*m/z*) 518 (*MH*<sup>+</sup>), *Rt* = 4.47, purity: 90 %.
- 1-[3-[[2-(5-Bromo -1*H*-indol-3-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12z**): LC/MS (*m/z*) 532 (*MH*<sup>+</sup>), *Rt* = 4.46, purity: 87 %.
- 15 1-[3-[[3-(5-Bromo -1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12aa**): LC/MS (*m/z*) 546 (*MH*<sup>+</sup>), *Rt* = 4.59, purity: 88 %.
- 1-[3-[[2-(5-Bromo -1*H*-indol-3-yl)ethyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12ab**): LC/MS (*m/z*) 546 (*MH*<sup>+</sup>), *Rt* = 4.50, purity: 90 %.
- 20 1-[3-[[4-(5-Bromo -1*H*-indol-3-yl)butyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12ac**): LC/MS (*m/z*) 560 (*MH*<sup>+</sup>), *Rt* = 4.61, purity: 90 %.
- 25 1-[3-[[3-(5-Bromo -1*H*-indol-3-yl)propyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12ad**): LC/MS (*m/z*) 560 (*MH*<sup>+</sup>), *Rt* = 4.62, purity: 92 %.
- 1-[3-[Ethyl[2-(5-iodo -1*H*-indol-3-yl)ethyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12ae**): LC/MS (*m/z*) 594 (*MH*<sup>+</sup>), *Rt* = 4.60, purity: 82 %.
- 30 %.

- 1-[3-[Ethyl[3-(5-iodo -1*H*-indol-3-yl)propyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12af**): LC/MS (*m/z*) 608 (*MH*<sup>+</sup>), *Rt* = 4.72, purity: 71 %.
- 5 1-[2-[[4-(5-Chloro -1*H*-indol-3-yl)butyl]methylamino]ethyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12ag**): LC/MS (*m/z*) 502 (*MH*<sup>+</sup>), *Rt* = 4.50, purity: 90 %.
- 1-[2-[[4-(5-Bromo -1*H*-indol-3-yl)butyl]methylamino]ethyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12ah**): LC/MS (*m/z*) 546 (*MH*<sup>+</sup>), *Rt* = 4.55, purity: 83 %.
- 10 1-[4-[[2-(5,7-Difluoro -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12ai**): LC/MS (*m/z*) 504 (*MH*<sup>+</sup>), *Rt* = 4.36, purity: 87 %.
- 1-[4-[[2-(7-Chloro -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12aj**): LC/MS (*m/z*) 502 (*MH*<sup>+</sup>), *Rt* = 4.42, purity: 70 %.
- 15 1-[4-[[2-(5-Chloro -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12ak**): LC/MS (*m/z*) 502 (*MH*<sup>+</sup>), *Rt* = 4.45, purity: 91 %.
- 1-[4-[[2-(5-Bromo -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12al**): LC/MS (*m/z*) 546 (*MH*<sup>+</sup>), *Rt* = 4.48, purity: 90 %.
- 20 1-[4-[[2-(5-Methyl -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12am**): LC/MS (*m/z*) 482 (*MH*<sup>+</sup>), *Rt* = 4.37, purity: 87 %.
- 25 1-[4-[[2-(5-Iodo -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12an**): LC/MS (*m/z*) 594 (*MH*<sup>+</sup>), *Rt* = 4.57, purity: 83 %.
- 1-[4-[[2-(5-(2-methyl-2-propyl)-1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12ao**): LC/MS (*m/z*) 524 (*MH*<sup>+</sup>), *Rt* = 4.85, purity: 91 %.
- 30

1-[4-[[2-(5-(2-Propyl)-1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**12ap**): LC/MS (*m/z*) 510 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 4.72, purity: 92 %.

5 **Example 13**

1-[3-[[2-(5-Methyl-1*H*-indol-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**13a**).

2-(3-Chloropropyl)-1,3-dioxolan-4-ylmethoxymethyl polystyrene (2.0 g, 1.6 mmol) was suspended in dry *N,N*-dimethylformamide (15 mL). Sodium iodide (0.67 g, 4.5 mmol) was added followed by diisopropylethylamine (1.70 mL, 9.6 mmol) and allyl amine (0.28 g, 4.8 mmol). The reaction mixture was heated at 80 °C under stirring for 12 h. After cooling to room temperature, the resin was filtered and washed with *N,N*-dimethylformamide (3 X 15 mL), methanol (3 X 15 mL), tetrahydrofuran (3 X 15 mL), and subsequently with methanol and tetrahydrofuran (each 10 mL, 5 cycles). Finally, the resin was washed with tetrahydrofuran (4 X 10 mL) and dried *in vacuo* (55 °C, 12 h). The resin was then suspended in dry *N,N*-dimethylformamide (20 mL). Sodium iodide (0.60 g, 4.0 mmol) was added followed by diisopropylethylamine (0.48 mL, 2.7 mmol) and 1-(3-chloropropyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**9**) (0.79 g, 2.5 mmol). The reaction mixture was stirred for 12 h at 80 °C. After cooling to room temperature, the resin was filtered and washed with *N,N*-dimethylformamide (3 X 15 mL), methanol (3 X 15 mL), tetrahydrofuran (3 X 15 mL), and then subsequently with methanol and tetrahydrofuran (each ca. 15 mL, 5 cycles). Finally, the resin was washed with tetrahydrofuran (4 X 15 mL) and dried *in vacuo* (55 °C, 12 h, 2.1 g).

An aliquot of this material (120 mg, ca. 0.08 mmol) and 4-methylphenylhydrazine hydrochloride (ca. 40 mg, 0.20 mmol) were mixed in a reactor tube. A 0.5 M solution of anhydrous zinc chloride in acetic acid (1.5 mL) was added and the reaction tube was sealed. The reaction mixture was stirred for 12 h at 75 °C. After cooling to room temperature, the reaction mixture was filtered and the residual resin washed with dimethylsulfoxide (1.5 mL). To the combined filtrates was added saturated aqueous sodium bicarbonate solution (1.5 mL). The solution was loaded on a reversed phase column (C-18, 1 g, Varian Mega Bond Elut®, Chrompack cat. no. 220508), pre-conditioned with methanol (3 mL) and water (3 mL). The column was washed with water

- (4 mL) and the product was eluted with methanol (4.5 mL). After evaporation of the volatile solvents, the crude product was purified by preparative reversed phase HPLC chromatography. The resulting solution was loaded on an ion exchange column (SCX, 1 g, Varian Mega Bond Elut®, Chrompack cat. no. 220776), pre-conditioned with 10 % solution of acetic acid in methanol (3 mL) and the column was washed with methanol (4 mL) and acetonitrile (4 mL), followed by elution with 4 N solution of ammonia in methanol (4.5 mL). Evaporation of the volatile solvents afforded the title compound (13a) as a colorless oil (2 mg, 4  $\mu$ mol, 5 %). LC/MS (m/z) 494 (MH<sup>+</sup>), Rt = 4.44, purity: 93 %.
- 10 The following compounds were prepared analogously:
- 1-[3-[[2-(5-Fluoro-1*H*-indol-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13b): LC/MS (m/z) 498 (MH<sup>+</sup>), Rt = 4.31, purity: 96 %.
- 1-[3-[[2-(7-Fluoro-1*H*-indol-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13c): LC/MS (m/z) 498 (MH<sup>+</sup>), Rt = 4.34, purity: 86 %.
- 1-[3-[[3-(5-Fluoro-1*H*-indol-3-yl)propyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13d): LC/MS (m/z) 512 (MH<sup>+</sup>), Rt = 4.48, purity: 96 %.
- 20 1-[3-[[3-(7-Fluoro-1*H*-indol-3-yl)propyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13e): LC/MS (m/z) 512 (MH<sup>+</sup>), Rt = 4.49, purity: 78 %.
- 1-[3-[[2-(5-Chloro-1*H*-indol-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13f): LC/MS (m/z) 514 (MH<sup>+</sup>), Rt = 4.52, purity: 86 %.
- 25 1-[3-[[2-(5,7-Difluoro-1*H*-indol-3-yl)ethyl]propylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13g): LC/MS (m/z) 518 (MH<sup>+</sup>), Rt = 4.47, purity: 89 %.
- 1-[3-[[2-[5-(2-Propyl)-1*H*-indol-3-yl]ethyl](2-propyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13h): LC/MS (m/z) 524 (MH<sup>+</sup>), Rt = 4.78, purity: 96 %.
- 30

- 1-[3-[[3-(4-Fluoro-7-methyl-1*H*-indol-3-yl)propyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13i): LC/MS (m/z) 526 (MH<sup>+</sup>), Rt = 4.65, purity: 83 %.
- 5 1-[3-[[2-(4-Chloro-7-methyl-1*H*-indol-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13j): LC/MS (m/z) 528 (MH<sup>+</sup>), Rt = 4.67, purity: 79 %.
- 1-[3-[[3-(5-Chloro-1*H*-indol-3-yl)propyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13k): LC/MS (m/z) 528 (MH<sup>+</sup>), Rt = 4.63, purity: 78 %.
- 10 1-[3-[[2-(5-Pyrrolo[3,2-*h*]-1*H*-quinolin-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13l): LC/MS (m/z) 531 (MH<sup>+</sup>), Rt = 3.43, purity: 91 %.
- 1-[3-[[3-(7-Fluoro-1*H*-indol-3-yl)propyl](2-furylmethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13m): LC/MS (m/z) 552 (MH<sup>+</sup>), Rt = 4.58, purity: 82 %.
- 15 1-[3-[[4-(7-Carboxy-1*H*-indol-3-yl)butyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13n): LC/MS (m/z) 552 (MH<sup>+</sup>), Rt = 4.17, purity: 69 %.
- 1-[3-[[2-[5-Bromo-1*H*-indol-3-yl]ethyl]-propylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13o): LC/MS (m/z) 560 (MH<sup>+</sup>), Rt = 4.62, purity: 96 %.
- 20 1-[3-[[3-(1*H*-Indol-3-yl)propyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13p): LC/MS (m/z) 574 (MH<sup>+</sup>), Rt = 4.78, purity: 93 %.
- 25 1-[3-[[2-(5-Methyl-1*H*-indol-3-yl)ethyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13q): LC/MS (m/z) 574 (MH<sup>+</sup>), Rt = 4.82, purity: 93 %.
- 1-[3-[[2-(5-Fluoro-1*H*-indol-3-yl)ethyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13r): LC/MS (m/z) 578 (MH<sup>+</sup>), Rt = 4.71, purity: 95 %.
- 30

1-[3-[[3-(1*H*-Pyrrolo[3,2-*h*]quinolin-3-yl)propyl](2-furylmethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13s): LC/MS (*m/z*) 585 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 3.60, purity: 90 %.

1-[3-[[3-(5-Methyl-1*H*-indol-3-yl)propyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13t): LC/MS (*m/z*) 588 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 4.96, purity: 82 %.

1-[3-[[3-(5-Fluoro-1*H*-indol-3-yl)propyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13u): LC/MS (*m/z*) 592 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 4.82, purity: 90 %.

1-[3-[[2-(5,7-Difluoro-1*H*-indol-3-yl)ethyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13v): LC/MS (*m/z*) 596 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 4.84, purity: 92 %.

1-[3-[[4-(1*H*-Pyrrolo[3,2-*h*]quinolin-3-yl)butyl](2-furylmethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13w): LC/MS (*m/z*) 599 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 3.71, purity: 83 %.

1-[3-[[2-(2-Phenoxyethyl)[2-[5-(2-propyl)-1*H*-indol-3-yl]ethyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13x): LC/MS (*m/z*) 602 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 5.24, purity: 78 %.

1-[3-[[2-(5-Bromo-1*H*-indol-3-yl)ethyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (13y): LC/MS (*m/z*) 638 (*MH*<sup>+</sup>), *R*<sub>t</sub> = 4.98, purity: 91 %.

#### Example 14

1-(3-Iodopropyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (14a). A solution/suspension of 1-(3-chloropropyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (20 g, 35 mmol, 80 % pure) and sodium iodide (285 g, 1.9 mol) in dry acetone (200 ml) was heated at reflux for 24 h. The mixture was evaporated, and partitioned between ether and water. The ether layer was separated, and was washed successively with water and brine. The organic extract was dried over anhydrous magnesium sulfate, filtered and evaporated to give 1-(3-iodopropyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (25,8 g, 99 %, 80 % pure) as a thick oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.6-1.9 (m, 2H),



2.21 (ddd, 1H), 2.31 (ddd, 1H), 3.16 (td, 2H), 5.12 (dt, 1H), 5.21 (dt, 1H), 7.02 (t, 2H), 7.41 (d, 2H), 7.43 (d, 1H), 7.51 (s, 1H), 7.62 (dq, 1H)

The following compounds were prepared analogously:

- 5 1-(2-Iodoethyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**14b**): yellow oil, <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 2.4-2.9 (m, 2H), 3.38 (dt, 1H), 3.46 (dt, 1H), 5.15 (d, 1H), 5.21 (d, 1H), 7.03 (t, 2H), 7.35-7.48 (m, 3H), 7.52 (s, 1H), 7.62 (d, 1H).

- 10 1-(4-Iodobutyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**14c**): yellow oil, <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.1-1.5 (m, 2H), 1.81 (tt, 2H), 2.00-2.30 (m, 2H), 3.11 (t, 2H), 5.14 (d, 1H), 5.20 (d, 1H), 7.01 (t, 2H), 7.35-7.47 (m, 3H), 7.51 (s, 1H), 7.60 (d, 1H).

#### Example 15

- 1-(3-(Ethylamino)propyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (**15a**). To a stirred solution of 1-(3-iodopropyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile (12.9 g, 30 mmol, 8 % pure) in ethanol (150 mL) was added a solution of ethylamine (20.3 g, 450 mmol) in THF (50 mL) portionwise, and the mixture was stirred over night. The solution was evaporated, and was dissolved/suspended in water. The pH was adjusted to 12 using aqueous sodium hydroxide solution (2 M) and was extracted with ether. The organic extract was washed with brine, dried over anhydrous magnesium sulfate, filtered and evaporated to give an oil. This oil was purified by silica chromatography using 50% v/v ethyl acetate/heptane as eluent, followed by 10% v/v triethylamine/ 40% v/v ethyl acetate/heptane followed by 20% v/v triethylamine/ethyl acetate to give the title compound (5.52 g, 57%) as a pale yellow oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.05 (t, 3 H), 1.2-1.6 (m, 2H), 2.15 (ddd, 1H), 2.24 (ddd, 1H), 2.57 (q, 2H) 2.58 (t, 2H), 5.12 (dt, 1H), 5.20 (dt, 1H), 7.00 (t, 2H), 7.38 (d, 1H), 7.42 (dd, 2H), 7.49 (s, 1H), 7.58 (ddt, 1H).

The following compounds were prepared analogously:

1-(2-(Methylamino)ethyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile

(15b): yellow oil;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.38 (s, 3H), 2.33-2.72 (m, 4H), 5.13 (d, 1H), 5.20 (d, 1H), 7.01 (t, 2H), 7.37-7.47 (m, 3H), 7.50 (s, 1H), 7.59 (d, 1H).

5 1-(4-(Methylamino)butyl)-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile

(15c): yellow oil;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.00-1.45 (m, 2H), 1.46 (tt, 2H), 2.10 (ddd, 1H), 2.21 (ddd, 1H), 2.37 (s, 3H), 2.50 (t, 2H), 5.13 (d, 1H), 5.19 (d, 1H), 7.00 (t, 2H), 7.34-7.46 (m, 3H), 7.49 (s, 1H), 7.59 (d, 1H).

10

### Pharmacological Testing

The affinity of the compounds of the invention to  $5\text{-HT}_{1A}$  receptors was determined by measuring the inhibition of binding of a radioactive ligand at  $5\text{-HT}_{1A}$  receptors as described

15 in the following test:

### Inhibition of $^3\text{H}$ -5-CT Binding to Human $5\text{-HT}_{1A}$ Receptors

By this method the inhibition by drugs of the binding of the  $5\text{-HT}_{1A}$  agonist

20  $^3\text{H}$ -5-carboxamido tryptamine ( $^3\text{H}$ -5-CT) to cloned human  $5\text{-HT}_{1A}$  receptors stably expressed in transfected HeLa cells (HA7) (Fargin, A. *et al*, *J. Biol. Chem.*, 1989, 264, 14848) is determined *in vitro*. The assay was performed as a modification of the method described by Harrington, M.A. *et al*, *J. Pharmacol. Exp. Ther.*, 1994, 268, 1098. Human  $5\text{-HT}_{1A}$  receptors (40  $\mu\text{g}$  of cell homogenate) were incubated for 15 minutes at 37 °C in 50

25 mM Tris buffer at pH 7.7 in the presence of  $^3\text{H}$ -5-CT. Non-specific binding was determined by including 10  $\mu\text{M}$  of metergoline. The reaction was terminated by rapid filtration through Unifilter GF/B filters on a Tomtec Cell Harvester. Filters were counted in a Packard Top Counter. The results obtained are presented in table 1 below.

30 The compounds of the invention have also been tested for their effect on re-uptake of serotonin in the following test:

### Inhibition of <sup>3</sup>H-5-HT Uptake Into Rat Brain Synaptosomes

Using this method, the ability of drugs to inhibit the accumulation of <sup>3</sup>H-5-HT into whole  
 5 rat brain synaptosomes is determined *in vitro*. The assay was performed as described by  
 Hyttel, J., *Psychopharmacology* 1978, 60, 13. The results obtained are presented in table 1:

Table 1

10

Compound No.	Inhibition of <sup>3</sup> H- 5-CT binding IC <sub>50</sub> ( nM) % inhibition at 100 nM	Inhibition of serotonin reuptake IC <sub>50</sub> ( nM) % inhibition at 100 nM
1a	39	60
1b	12	13
1c	53	85
2a	1.0	340
2b	6.4	40
2e	38	15
2f	8.6	14
2g	40	20
2j	41	9.7
2m	4.7	Not tested
2n	15	Not tested
2o	12	31
4a	23	54
4b	63	59% inh. at 100 nM
4c	11	4% inh. at 100 nM
4d	4.5	7% inh. at 100 nM

4e	17	160
4f	1.6	4% inh. at 100 nM
4g	18	28% inh. at 100 nM
4h	3.2	69
4i	1.9	26% inh. at 100 nM
4j	6.1	78
4k	0.42	100
4l	76% inh. at 100 nM	27% inh. at 100 nM
4m	65% inh. at 100 nM	74% inh. at 100 nM
4n	14	39% inh. at 100 nM
4o	26	73
4p	19	6% inh. at 100 nM
4q	16	60% inh. at 100 nM
4r	11	19% inh. at 100 nM
4s	30	35
4t	69% inh. at 100 nM	73% inh. at 100 nM
4u	58% inh. at 100 nM	44% inh. at 100 nM
12b	43	10
12c	19	17
12d	31	12
12f	4.7	13
12i	27	20
12j	7.9	14
12k	3.6	8.4
12o	6.2	49% inh. at 100 nM
12p	19	11

12q	12	6.3
12r	16	47% inh. at 100 nM
12s	7.7	18
12u	9.0	22
12v	39	12
12x	14	50% inh. at 100 nM
12aa	16	37% inh. at 100 nM
12ab	20	50% inh. at 100 nM
12ad	21	35% inh. at 100 nM
12ae	11	49% inh. at 100 nM
12af	31	38% inh. at 100 nM
13b	7.4	44
13c	9.6	12
13d	15	21
13e	22	27
13f	31	16% inh. at 100 nM
13g	18	49% inh. at 100 nM
13j	16	61% inh. at 100 nM
13k	19	Not tested
13p	23	Not tested
13q	12	Not tested
13r	8.9	Not tested
13t	23	Not tested
13u	22	Not tested
13v	23	Not tested
13x	26	Not tested
Pindolol	100	
Paroxetine	-	0.29

Table 1 reference compounds

Furthermore, the 5-HT<sub>1A</sub> antagonistic activity of some of the compounds of the invention has been estimated *in vitro* at cloned 5-HT<sub>1A</sub> receptors stably expressed in transfected HeLa cells (HA7). In this test, 5-HT<sub>1A</sub> antagonistic activity is estimated by measuring the ability of the compounds to antagonize the 5-HT induced inhibition of forskolin induced cAMP accumulation. The assay was performed as a modification of the method described by Pauwels, P.J. *et al*, *Biochem. Pharmacol.* 1993, 45, 375.

As seen from the above, the compounds of the invention show affinity for the 5-HT<sub>1A</sub> receptor. Furthermore, many of the compounds of the present invention possess valuable activity as serotonin re-uptake inhibitors.

Accordingly, the compounds are considered useful for the treatment of psychiatric and neurological disorders as mentioned previously.

#### Pharmaceutical formulation

The pharmaceutical formulations of the invention may be prepared by conventional methods in the art. For example: Tablets may be prepared by mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvants or additives usually used for such purposes such as colourings, flavourings, preservatives etc. may be used provided that they are compatible with the active ingredients.

Solutions for injections may be prepared by dissolving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to desired volume, sterilisation of the solution and filling in suitable ampules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.


The pharmaceutical compositions of this invention or those which are manufactured in accordance with this invention may be administered by any suitable route, for example orally in the form of tablets, capsules, powders, syrups, etc., or parenterally in the form of

solutions for injection. For preparing such compositions, methods well known in the art may be used, and any pharmaceutically acceptable carriers, diluents, excipients, or other additives normally used in the art may be used.

Conveniently, the compounds of the invention are administered in unit dosage form  
5 containing said compounds in an amount of about 0.01 to 1000 mg. The total daily dose is usually in the range of about 0.05 - 500 mg, and most preferably about 0.1 to 50 mg of the active compound of the invention.

**Claims:**

1. An isobenzofuran having the general Formula I:







wherein

- 10 R¹ is hydrogen, halogen, trifluoromethyl, trifluoromethylsulfonyloxy, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-8</sub> cycloalkyl, C<sub>1-6</sub> alkoxy, hydroxy, formyl, acyl, amino, C<sub>1-6</sub> alkylamino, C<sub>2-12</sub> dialkylamino, acylamino, C<sub>1-6</sub> alkoxycarbonylamino, aminocarbonylamino, C<sub>1-6</sub> alkylaminocarbonylamino, C<sub>2-12</sub> dialkylaminocarbonylamino, 15 nitro, cyano, COOH, or COO-C<sub>1-6</sub> alkyl;

R² and R³ are each independently selected from hydrogen, trifluoromethyl, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-8</sub> cycloalkyl and C<sub>1-6</sub> alkoxy;

- 20 n is 1, 2, 3, 4 or 5;  
m is 0 or 1;

A is selected from the following groups:



- 25 wherein  
Z is O or S;  
s is 0 or 1;




q is 0 or 1;

R<sup>4</sup> is hydrogen, C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl, C<sub>2-6</sub>-alkynyl, C<sub>1-6</sub>-alkyl-Aryl, or C<sub>1-6</sub>-alkyl-O-Aryl;

- 5 D is a spacer group selected from branched or straight chain C<sub>1-6</sub>-alkylene, C<sub>2-6</sub>-alkenylene and C<sub>2-6</sub>-alkynylene;

B is a group selected from a group of formula (II), (III), and (IV)



- 10 wherein R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup> and R<sup>10</sup> are each independently selected among the R<sup>1</sup> substituents;

or R<sup>8</sup> and R<sup>9</sup> together form a fused 5- or 6-membered ring optionally containing further heteroatoms; and the resulting heterocycle is optionally substituted with substituents

- 15 selected among the R<sup>1</sup> substituents;

or two of the groups of R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are linked together thereby forming a

—O—(CH<sub>2</sub>)<sub>p</sub>—O— -bridge wherein p is 1 or 2;

- 20 Ar and Aryl are independently selected from the group consisting of phenyl, 2-thienyl, 3-thienyl, 2-furanyl, 3-furanyl, 2-pyrimidyl, 1-indolyl, 2-indolyl, 3-indolyl, 1-indol-2-onyl, 3-indol-2-onyl, 2- or 3-benzofuranyl, 2- or 3-benzothiophenyl, 1-naphthyl or 2-naphthyl, each optionally substituted with halogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylthio, hydroxy, C<sub>1-6</sub> alkylsulfonyl, cyano, trifluoromethyl, trifluoromethylsulfonyloxy, C<sub>3-8</sub>
- 25 cycloalkyl, C<sub>3-8</sub> cycloalkyl-C<sub>1-6</sub> alkyl, nitro, amino, C<sub>1-6</sub> alkylamino, C<sub>2-12</sub> dialkylamino, acylamino or alkylenedioxy;

its enantiomers, and pharmaceutically acceptable acid addition salt thereof.

2. A compound of Claim 1, **characterised in** that A is a group of formula (1).
3. A compound of Claim 1, **characterised in** that A is a group of formula (2).
- 5 4. A compound of Claim 1, **characterised in** that A is a group of formula (3).
5. A compound of Claim 1, **characterised in** that A is a group of formula (4).
- 10 6. A compound of Claim 2, **characterised in** that R<sup>4</sup> is methyl, ethyl, propyl, 2-propen-1-yl, 2-furylmethyl, 2-phenoxyethyl;
7. A compound of any of the Claims 2 and 6, **characterised in** that q = 0;
- 15 8. A compound of any of the Claims 2 and 6, **characterised in** that q = 1 and Z is O.
9. A compound of any of the Claims 1 - 8, **characterised in** that B is a group of formula (II).
- 20 10. A compound of any of the Claims 1 - 8, **characterised in** that B is a group of formula (III).
11. A compound of any of the Claims 1 - 8, **characterised in** that B is a group of formula (IV).
- 25 12. A compound of Claim 9, **characterised in** that at least one of R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup>, is methoxy.
13. A compound of Claim 9, **characterised in** that Formula (II) is a benzodioxan group
- 30 or a 1,2-methylenedioxybenzene group.
14. A compound of Claim 10, **characterised in** that Formula (III) is a 3-indolyl.

15. A compound of Claim 14, **characterised in** that the 3-indolyl is substituted in 5-position by methyl, fluoro, chloro, bromo, iodo, *t*-butyl or *i*-propyl, or in 7-position by fluoro, chloro or carboxy; or disubstituted by 5,7-difluoro, 4-fluoro-7-methyl or 4-chloro-  
5 7-methyl or the two substituents together form a pyridyl ring fused to the 3-indolyl.
16. A compound of Claim 11, **characterised in** that Formula (IV) is a 4-indolyl or a 5-indolyl group.
- 10 17. A compound of any of the Claims 1 - 16, **characterised in** that Ar is phenyl or phenyl substituted with halogen or CF<sub>3</sub>;
18. A compound of Claim 17, **characterised in** that Ar is phenyl which may be substituted with Cl or F in the 4-position or Cl or CF<sub>3</sub> in the 3-position.
- 15 19. A compound of any of the Claims 1 - 18, **characterised in** that R<sup>1</sup> is H, CN or F in the 5-position of the isobenzofuran group.
- 20 20. A compound of any of the Claims 1 - 19, **characterised in** that R<sup>2</sup> and R<sup>3</sup> are selected from hydrogen or methyl.
21. A compound of any of the Claims 1 - 20, **characterised in** that n = 2, 3 or 4.
22. A compound of claim 21, **characterised in** that n = 3;
- 25 23. A compound of any of the Claims 1 - 22, **characterised in** that m = 0.
24. A compound of any of the Claims 1, 18, 19, 20, 21, 22 and 23, **characterised in** that R<sup>2</sup> and R<sup>3</sup> are both hydrogen; R<sup>1</sup> is H, CN or F in the 5-position of the isobenzofuran  
30 group; and Ar is phenyl which may be substituted with F or Cl in the 4-position or with Cl or CF<sub>3</sub> in the 3-position.

25. A compound of any of the Claims 1 and 24, **characterised in** that A is a group of formula (1);  $q = 0$ ;  $R^4$  is methyl; D is propylene;  $m = 0$ ; and B is a 1,4-benzodioxan group of Formula (II) attached in the 5-position.
- 5 26. A compound of any of the Claims 1 and 24, **characterised in** that A is a group of formula (1);  $R^4$  is  $CH_3$  or prop-2-en-1-yl;  $n = 3$ ; D is ethylene or propylene; and B is a phenyl group wherein at least one substituent is OMe.
- 10 27. A compound of any of the Claims 1 and 24, **characterised in** that A is a group of formula (1);  $q$  is 0;  $R^4$  is methyl, ethyl, propyl, 2-propen-1-yl, 2-furylmethyl or 2-phenoxyethyl; D is ethylene or propylene;  $m = 0$ ; and B is a 3-indolyl group of Formula (III).
- 15 28. A compound according to claim 27, **characterised in** that the 3-indolyl group is substituted by methyl, fluoro, chloro, bromo, iodo, *t*-butyl or *i*-propyl in the 5-position; or fluoro, chloro or carboxy in the 7-position; or by 5,7-difluoro, 4-fluoro-7-methyl or 4-chloro-7-methyl; or the two substituents together form a pyridyl ring fused to the 3-indolyl-group.
- 20 29. A compound of any of the Claims 1 and 24, **characterised in** that A is a group of formula (2) or (3);  $n = 3$ ;  $m = 0$ ; and B is an 4- or 5-indolyl-group of Formula (IV) wherein  $R^{10}$  is hydrogen;  $R^1$  is CN in the 5-position of the isobenzofuran and Ar is 4-Fluorophenyl.
30. The compound according to claim 1 which is
- 25 (-)-1-[3-[[4-(1,4-Benzodioxan-5-yl)butyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(1,4-Benzodioxan-5-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile oxalate,
- 1-[3-[[2-(1,4-Benzodioxan-5-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-
- 30 dihydroisobenzofuran-5-carbonitrile oxalate,

- 1-[3-[[1,4-Benzodioxan-5-ylmethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile oxalate,
- 1-(4-Fluorophenyl)-1-[3-[4-(2-methoxyphenyl)piperazinyl]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,
- 5 1-(4-Fluorophenyl)-1-[3-[methyl[2-(2-methoxyphenoxy)ethyl]amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-(4-Fluorophenyl)-1-[3-[methyl[2-(3-methoxyphenoxy)ethyl]amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,
- (S)-1-[3-[[4-(1*H*-Indol-3-yl)butyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 10 1-[3-[[4-(1*H*-Indol-3-yl)butyl]methylamino]propyl]-1-phenyl-1,3-dihydroisobenzofuran, (S)-1-[3-[[3-(1*H*-Indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(1*H*-Indol-3-yl)propyl]methylamino]propyl]-1-phenyl-1,3-dihydroisobenzofuran,
- 15 5-[3-[[3-(1-Phenyl-1,3-dihydroisobenzofuran-1-yl)propyl]methylamino]propyl]-1,4-benzodioxane,
- 5-[3-[[3-[1-(3-Chlorophenyl)-1,3-dihydroisobenzofuran-1-yl]propyl]methylamino]propyl]-1,4-benzodioxane,
- 5-[3-[[3-[1-(4-Fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]propyl]methylamino]propyl]-1,4-benzodioxane,
- 20 5-[3-[[3-[1-(3-Trifluoromethylphenyl)-1,3-dihydroisobenzofuran-1-yl]propyl]methylamino]propyl]-1,4-benzodioxane,
- 1-[3-[[3-(1,4-Benzodioxan-5-yl)propyl]methylamino]propyl]-1-(4-chlorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 25 1-[3-[4-(1*H*-Indol-4-yl)piperazinyl]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[4-(1*H*-Indol-5-yl)piperazinyl]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[4-(1*H*-Indol-3-yl)piperidinyl]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-
- 30 5-carbonitrile,

- 5-[3-[[3-[-5-Fluoro-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]propyl]methylamino]propyl]-1,4-benzodioxane,  
 1-[3-[[2-(1*H*-Indolyl-3-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,  
 5 1-(4-Fluorophenyl)-1-[3-[[2-(3-methoxyphenyl)ethyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,  
 1-(4-Fluorophenyl)-1-[3-[[2-(3-methoxyphenyl)ethyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,  
 1-(4-Fluorophenyl)-1-[3-[[2-(2-methoxyphenyl)ethyl](prop-2-en-1-yl)amino]propyl]-1,3-  
 10 dihydroisobenzofuran-5-carbonitrile,  
 1-[3-[[2-(2,5-Dimethoxyphenyl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,  
 1-[3-[[2-(2,5-Dimethoxyphenyl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,  
 15 1-(4-Fluorophenyl)-1-[3-[[2-phenoxyethyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,  
 1-[3-[[2-(1*H*-Indolyl-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,  
 1-(4-Fluorophenyl)-1-[3-[[2-phenoxyethyl](prop-2-en-1-yl)amino]propyl]-1,3-  
 20 dihydroisobenzofuran-5-carbonitrile,  
 1-(4-Fluorophenyl)-1-[3-[[3-(2-methoxyphenyl)propyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,  
 1-(4-Fluorophenyl)-1-[3-[[3-(2-methoxyphenyl)propyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,  
 25 1-(4-Fluorophenyl)-1-[3-[[3-(3-methoxyphenyl)propyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,  
 1-(4-Fluorophenyl)-1-[3-[[3-(2-methoxyphenoxy)propyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,  
 1-(4-Fluorophenyl)-1-[3-[[3-(2-methoxyphenoxy)propyl](prop-2-en-1-yl)amino]propyl]-  
 30 1,3-dihydroisobenzofuran-5-carbonitrile,

- 1-(4-Fluorophenyl)-1-[3-[[3-(3-methoxyphenoxy)propyl]methylamino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-(4-Fluorophenyl)-1-[3-[[3-(3-methoxyphenoxy)propyl](prop-2-en-1-yl)amino]propyl]-1,3-dihydroisobenzofuran-5-carbonitrile,
- 5 1-[3-[(2-Benzyloxyethyl)methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[(2-Benzyloxyethyl)(prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(1*H*-Indolyl-3-yl)propyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 10 1-[3-[[3-(1*H*-Indolyl-3-yl)propyl](2-propynyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(1*H*-Indolyl-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 15 1-[3-[[2-(5-Methyl-1*H*-indol-3-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(7-Fluoro-1*H*-indol-3-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 5-Fluoro-1-[3-[[3-(5-methyl-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran,
- 20 5-Fluoro-1-[3-[[3-(7-fluoro-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran,
- 1-[3-[[3-(5-Methyl-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 25 1-[3-[Ethyl[3-(1*H*-indol-3-yl)propyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[Ethyl[2-(5-methyl-1*H*-indol-3-yl)ethyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(7-Fluoro-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 30

- 1-[3-[[3-(5-Fluoro-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[Ethyl[2-(5-fluoro-1*H*-indol-3-yl)ethyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 5 1-[3-[Ethyl[2-(7-fluoro-1*H*-indol-3-yl)ethyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(5-Chloro-1*H*-indol-3-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(5-Chloro-1*H*-indol-3-yl)propyl]methylamino]propyl]-5-fluoro-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran,
- 10 1-[3-[[4-(5-Methyl-1*H*-indol-3-yl)butyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[Ethyl[3-(5-methyl-1*H*-indol-3-yl)propyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 15 1-[3-[Ethyl[3-(7-fluoro-1*H*-indol-3-yl)propyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[Ethyl[3-(5-fluoro-1*H*-indol-3-yl)propyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(5-Chloro-1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 20 1-[3-[[2-(7-Chloro-1*H*-indol-3-yl)ethyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(5-Chloro-1*H*-indol-3-yl)ethyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 25 1-[3-[[2-(5,7-Difluoro-1*H*-indol-3-yl)ethyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[4-(5-Fluoro-1*H*-indol-3-yl)butyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[4-(5-Chloro-1*H*-indol-3-yl)butyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 30 1-[3-[[3-(5-Chloro-1*H*-indol-3-yl)propyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,



- 1-[3-[[3-(5,7-Difluoro -1*H*-indol-3-yl)propyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(5-Bromo -1*H*-indol-3-yl)ethyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 5 1-[3-[[3-(5-Bromo -1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(5-Bromo -1*H*-indol-3-yl)ethyl]ethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[4-(5-Bromo -1*H*-indol-3-yl)butyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 10 1-[3-[[3-(5-Bromo -1*H*-indol-3-yl)propyl]methylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[Ethyl[2-(5-iodo -1*H*-indol-3-yl)ethyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 15 1-[3-[Ethyl[3-(5-iodo -1*H*-indol-3-yl)propyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[2-[[4-(5-Chloro -1*H*-indol-3-yl)butyl]methylamino]ethyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[2-[[4-(5-Bromo -1*H*-indol-3-yl)butyl]methylamino]ethyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 20 1-[4-[[2-(5,7-Difluoro -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[4-[[2-(7-Chloro -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 25 1-[4-[[2-(5-Chloro -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[4-[[2-(5-Bromo -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[4-[[2-(5-Methyl -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 30 1-[4-[[2-(5-Iodo -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,

- 1-[4-[[2-(5-*t*-Butyl -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[4-[[2-(5-*i*-Propyl -1*H*-indol-3-yl)ethyl]methylamino]butyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 5 1-[3-[[2-(5-Methyl -1*H*-indol-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(5-Fluoro -1*H*-indol-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(7-Fluoro -1*H*-indol-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 10 1-[3-[[3-(5-Fluoro -1*H*-indol-3-yl)propyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(7-Fluoro -1*H*-indol-3-yl)propyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 15 1-[3-[[2-(5-Chloro -1*H*-indol-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(5,7-Difluoro -1*H*-indol-3-yl)ethyl]-propylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-[5-(2-Propyl)-1*H*-indol-3-yl]ethyl]-2-propylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 20 1-[3-[[3-(4-Fluoro-7-methyl-1*H*-indol-3-yl)propyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(4-Chloro-7-methyl-1*H*-indol-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 25 1-[3-[[3-(5-Chloro -1*H*-indol-3-yl)propyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(5-Pyrrolo[3,2-*h*]-1*H*-quinolin-3-yl)ethyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(7-Fluoro -1*H*-indol-3-yl)propyl](2-furylmethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 30 1-[3-[[4-(7-Carboxy-1*H*-indol-3-yl)butyl](prop-2-en-1-yl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,

- 1-[3-[[2-[5-Bromo-1*H*-indol-3-yl]ethyl]-propylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(1*H*-Indol-3-yl)propyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 5 1-[3-[[2-(5-Methyl-1*H*-indol-3-yl)ethyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[2-(5-Fluoro-1*H*-indol-3-yl)ethyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(5-Pyrrolo[3,2-*h*]-1*H*-quinolin-3-yl)propyl]-2-furylmethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 10 1-[3-[[3-(5-Methyl-1*H*-indol-3-yl)propyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[3-(5-Fluoro-1*H*-indol-3-yl)propyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 15 1-[3-[[2-(5,7-Difluoro-1*H*-indol-3-yl)ethyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[[4-(5-Pyrrolo[3,2-*h*]-1*H*-quinolin-3-yl)butyl]-2-furylmethylamino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile,
- 1-[3-[2-Phenoxyethyl[2-[5-(2-propyl)-1*H*-indol-3-yl]ethyl]amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile or
- 20 1-[3-[[2-(5-Bromo-1*H*-indol-3-yl)ethyl](2-phenoxyethyl)amino]propyl]-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-5-carbonitrile
- or an acid addition salt thereof.
- 25 31. A pharmaceutical composition comprising a compound according to claims 1 to 30 or a pharmaceutically acceptable acid addition salt thereof and at least one pharmaceutically acceptable carrier or diluent.
32. The use of a compound according to claims 1 to 30 or a pharmaceutically
- 30 acceptable acid addition salt thereof for the preparation of a medicament for the treatment of a disorder or disease responsive to the effect of 5-HT<sub>1A</sub> receptors.

33. The use of a compound according to claim 32 wherein the medicament is for the treatment of depression, psychosis, anxiety disorders, panic disorder, obsessive compulsive disorder, impulse control disorder, alcohol abuse, aggression, ischaemia, senile dementia, cardiovascular disorders and social phobia.

5

34. A method for the treatment of a disorder or disease of living animal body, including a human, which is responsive to the effect of 5-HT<sub>1A</sub> receptors comprising administering to such a living animal body, including a human, a therapeutically effective amount of a compound according to claims 1 to 30 or a pharmaceutically acceptable acid addition salt thereof.

10

35. A method of treatment according to claim 34 where the disorder or disease is depression, psychosis, anxiety disorders, panic disorder, obsessive compulsive disorder, impulse control disorder, alcohol abuse, aggression, ischaemia, senile dementia, cardiovascular disorders or social phobia.

15

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00676

## A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 307/87, A61K 31/343

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 2657013 A1 (KEFALAS A/S), 28 July 1977 (28.07.77)  --	1-32
A	GB 1173312 A (KEFALAS A/S), 10 December 1969 (10.12.69)  --	1-32
A	WO 9518118 A1 (THE UPJOHN COMPANY), 6 July 1995 (06.07.95)  -----	1-32

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

29 March 2000

Date of mailing of the international search report

17-04-2000

Name and mailing address of the ISA/  
 Swedish Patent Office  
 Box 5055, S-102 42 STOCKHOLM  
 Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson/EÖ  
 Telephone No. +46 8 782 25 00

# INTERNATIONAL SEARCH REPORT

International application No.  
**PCT/DK 99/00676**

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **34-35**  
because they relate to subject matter not required to be searched by this Authority, namely:  
**A method for treatment of the human or animal body by therapy, see rule 39.1.**
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT  
Information on patent family members

02/12/99

International application No.  
PCT/DK 99/00676

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2657013 A1	28/07/77	AT 359488 B	10/11/80
		AT 360001 B	10/12/80
		AT 360002 B	10/12/80
		AT 571979 A	15/05/80
		AT 572079 A	15/05/80
		AT 947276 A	15/04/80
		AU 509445 B	15/05/80
		AU 2107377 A	13/07/78
		BE 850401 A	14/07/77
		CA 1094087 A	20/01/81
		CH 626886 A	15/12/81
		CH 632258 A	30/09/82
		CH 632259 A	30/09/82
		DK 13177 A	15/07/77
		ES 454980 A	01/04/78
		FI 63754 B,C	29/04/83
		FI 770073 A	15/07/77
		FR 2338271 A,B	12/08/77
		GB 1526331 A	27/09/78
		IE 44055 B	29/07/81
		JP 1368581 C	11/03/87
		JP 52105162 A	03/09/77
		JP 61035986 B	15/08/86
		NL 192451 B,C	01/04/97
		NL 7700244 A	18/07/77
		NO 147243 B	22/11/82
		NO 770109 A	15/07/77
		NZ 183001 A	02/06/78
		SE 429551 B,C	12/09/83
		SE 7614201 A	15/07/77
		US 4136193 A	23/01/79
		ZA 7700057 A	30/11/77
GB 1173312 A	10/12/69	BE 695302 A	14/08/67
		CH 495975 A	15/09/70
		CH 511828 A	31/08/71
		DE 1593918 A	27/04/72
		DK 120640 B	28/06/71
		DK 120761 B	12/07/71
		DK 120762 B	12/07/71
		DK 120956 B	09/08/71
		DK 121026 B	23/08/71
		FR 1514754 A	00/00/00
		NL 6703332 A	18/09/67
		NO 121896 B	26/04/71
		SE 354862 B	26/03/73
		US 3523124 A	04/08/70
		US 3629444 A	21/12/71

# INTERNATIONAL SEARCH REPORT

Information on patent family members

02/12/99

International application No.

PCT/DK 99/00676

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9518118 A1	06/07/95	AT 182590 T	15/08/99
		AU 684874 B	08/01/98
		AU 1257195 A	17/07/95
		CA 2176722 A	06/07/95
		CN 1139431 A	01/01/97
		DE 69419778 D	00/00/00
		EP 0737189 A,B	16/10/96
		SE 0737189 T3	
		ES 2133714 T	16/09/99
		IL 111940 D	00/00/00
		JP 9507224 T	22/07/97
		NZ 277373 A	28/07/98
		US 5877317 A	02/03/99
		ZA 9409927 A	13/06/96
-----			